



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N	A2	(11) International Publication Number: WO 95/34634
		(43) International Publication Date: 21 December 1995 (21.12.95)
<p>(21) International Application Number: PCT/EP95/02157</p> <p>(22) International Filing Date: 6 June 1995 (06.06.95)</p> <p>(30) Priority Data: 08/254,776 6 June 1994 (06.06.94) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 08/254,776 (CIP) Filed on 6 June 1994 (06.06.94)</p> <p>(71) Applicant (for all designated States except US): PLANT GENETIC SYSTEMS, N.V. [BE/BE]; Kolonel Bourgstraat 106, B-1040 Brussels (BE).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): KREBBERS, Enno [US/US]; 2014 Branard, Houston, TX 77098 (US). WILLIAMS, Mark [US/BE]; Holdaal 31, B-9000 Gent (BE). LEEMANS, Jan [BE/BE]; Wijngaard 7, B-9831 Deurle (BE).</p> <p>(74) Agents: GUTMANN, Ernest et al.; E. Gutmann - Y. Plasseraud S.A., 3, rue Chauveau-Lagarde, F-75008 Paris (FR).</p>	<p>(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>	
(54) Title: USE OF ANTHOCYANIN GENES TO MAINTAIN MALE STERILE PLANTS		
<p>(57) Abstract</p> <p>A plant consisting essentially of cells which comprise in their genome a homozygous male-sterility genotype at a first genetic locus; and a color-linked restorer genotype at a second genetic locus, which is heterozygous (Rf/-) for a foreign DNA Rf. The foreign DNA Rf comprises: a) a fertility-restorer gene capable of preventing the phenotypic expression of the male-sterility genotype, and b) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of the plant which is capable of producing anthocyanin at least in the seeds of the plant, so that anthocyanin production in the seeds is visible externally. Preferably, the anthocyanin regulatory gene is a shortened R, B or C1 gene or a continuation thereof. The invention also relates to DNA sequences encoding shortened R, B or C1 anthocyanin regulatory genes and to a process for maintaining a line of male-sterile plants which comprises crossing a male-sterile parent plant and a maintainer parent plant comprising homozygous male-sterility genotype and a restore genotype comprising fertility-restorer gene and an anthocyanin regulatory gene.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

USE OF ANTHOCYANIN GENES TO MAINTAIN MALE STERILE PLANTS

The present invention relates to a method to maintain male-sterile plants that can be used for the production of hybrid seed of a plant crop species, to transgenic inbred plants that can be used in such process, and to chimeric genes that can be used to produce such transgenic inbred plants.

Background to the Invention

In many, if not most plant species, the development of hybrid cultivars is highly desired because of their generally increased productivity due to heterosis: the superiority of performance of hybrid individuals compared with their parents (see e.g. Fehr, 1987, Principles of cultivar development, Volume 1 : Theory and Technique, MacMillan Publishing Company, New York; Allard, 1960, Principles of Plant Breeding, John Wiley and Sons, Inc.).

The development of hybrid cultivars of various plant species depends upon the capability of achieving essentially almost complete cross-pollination between parents. This is most simply achieved by rendering one of the parent lines male sterile (i.e. bringing them in a condition so that pollen is absent or nonfunctional) either manually, by removing the anthers, or genetically

by using, in the one parent, cytoplasmic or nuclear genes that prevent anther and/or pollen development (for a review of the genetics of male sterility in plants see Kaul, 1988, 'Male Sterility in Higher Plants', Springer Verlag).

For hybrid plants where the seed is the harvested product (e.g. corn, oilseed rape) it is in most cases also necessary to ensure that fertility of the hybrid plants is fully restored. In systems in which the male sterility is under genetic control this requires the existence and use of genes that can restore male fertility. The development of hybrid cultivars is mainly dependent on the availability of suitable and effective sterility and restorer genes.

Endogenous nuclear loci are known for most plant species that may contain genotypes which effect male sterility, and generally, such loci need to be homozygous for particular recessive alleles in order to result in a male-sterile phenotype. The presence of a dominant 'male fertile' allele at such loci results in male fertility.

Recently it has been shown that male sterility can be induced in a plant by providing the genome of the plant with a chimeric male-sterility gene comprising a DNA sequence (or male-sterility DNA) coding, for example, for a cytotoxic product (such as an RNase) and under the control of a promoter which is predominantly active in selected tissue of the male reproductive organs. In this

regard stamen-specific promoters, such as the promoter of the TA29 gene of Nicotiana tabacum, have been shown to be particularly useful for this purpose (Mariani et al., 1990, Nature 347:737, European patent publication ("EP") 0,344,029). By providing the nuclear genome of the plant with such a male-sterility gene, an artificial male-sterility locus is created containing the artificial male-sterility genotype that results in a male-sterile plant.

10 In addition it has been shown that male fertility can be restored to the plant with a chimeric fertility-restorer gene comprising another DNA sequence (or fertility-restorer DNA) that codes, for example, for a protein that inhibits the activity of the cytotoxic product or otherwise prevents the cytotoxic product from being active in the plant cells (European patent publication "EP" 0,412,911). For example the barnase gene of Bacillus amyloliquefaciens codes for an RNase, called barnase, which can be inhibited by a protein, barstar, that is encoded by the barstar gene of B. amyloliquefaciens. The barnase gene can be used for the construction of a sterility gene while the barstar gene can be used for the construction of a fertility-restorer gene. Experiments in different plant species, e.g. oilseed rape, have shown that a chimeric barstar gene can fully restore the male fertility of male sterile lines in which the male sterility was due to the presence of a

chimeric barnase gene (EP 0,412,911, Mariani et al., 1991, Proceedings of the CCIRC Rapeseed Congress, July 9-11, 1991, Saskatoon, Saskatchewan, Canada; Mariani et al., 1992, Nature 357:384). By coupling a marker gene, such as a dominant herbicide resistance gene (for example the bar gene coding for phosphinothricin acetyl transferase (PAT) that converts the herbicidal phosphinothricin to a non-toxic compound [De Block et al., 1987, EMBO J. 6:2513]), to the chimeric male-sterility and/or fertility-restorer gene, breeding systems can be implemented to select for uniform populations of male sterile plants (EP 0,344,029; EP 0,412,911).

The production of hybrid seed of any particular cultivar of a plant species requires the: 1) maintenance of small quantities of pure seed of each inbred parent, and 2) the preparation of larger quantities of seed of each inbred parent. Such larger quantities of seed would normally be obtained by several (usually two) seed multiplication rounds, starting from a small quantity of pure seed ("basic seed") and leading, in each multiplication round, to a larger quantity of pure seed of the inbred parent and then finally to a stock of seed of the inbred parent (the "parent seed" or "foundation seed") which is of sufficient quantity to be planted to produce the desired quantities of hybrid seed. Of course,

in each seed multiplication round larger planting areas (fields) are required.

In order to maintain and enlarge a small stock of seeds that can give rise to male-sterile plants it is necessary to cross the male sterile plants with normal pollen-producing parent plants. In the case in which the male-sterility is encoded in the nuclear genome, the offspring of such cross will in all cases be a mixture of male-sterile and male-fertile plants and the latter have to be removed from the former. With male-sterile plants containing an artificial male-sterility locus as described above, such removal can be facilitated by genetically linking the chimeric male sterility gene to a suitable marker gene, such as the bar gene, which allows the easy identification and removal of male-fertile plants (e.g. by spraying of an appropriate herbicide).

However, even when suitable marker genes are linked to male-sterility genotypes, the maintenance of parent male-sterile plants still requires at each generation the removal from the field of a substantial number of plants. For instance in systems using a herbicide resistance gene (e.g. the bar gene) linked to a chimeric male-sterility gene, as outlined above, only half of the parent stock will result in male-sterile plants, thus requiring the removal of the male-fertile plants by herbicide spraying prior to flowering. In any given

field, the removal of male-fertile plants effectively reduces the potential yield of hybrid seed or the potential yield of male-sterile plants during each round of seed multiplication for producing parent seed. In addition removal of the male- fertile plants may lead to irregular stands of the male-sterile plants. For these reasons removal of the male-fertile plants is economically unattractive for many important crop species such as corn and oilseed rape.

Anthocyanins are pigments that are responsible for many of the red and blue colors in plants. The genetic basis of anthocyanin biosynthesis has been well characterized, particularly in corn, Petunia, and Antirrhinium (Dooner et al, 1991, Ann.Rev.Genet. 25:179-199; Jayaram and Peterson, 1990, Plant Breeding Reviews 2:91-137; Coe, 1994, In 'The Maize Handbook', Freeling and Walbot, eds. Springer Verlag New York Inc., p. 279-281). In corn anthocyanin biosynthesis is apparently under control of 20 or more genes. The structural loci C2, Whp, A1, A2, Bz1, and Bz2 code for various enzymes involved in anthocyanin biosynthesis and at least 6 regulatory loci, acting upon the structural genes, have been identified in corn i.e. the R, B, C1, P1, P and Vp1 loci.

The R locus has turned out to be a gene family (in corn located on chromosome 10) comprising at least three

different genes i.e. R (which itself may comprise duplicate genes organized in a tandem array), and the displaced duplicate genes R(Sn) and R(Lc). R typically conditions pigmentation of the aleurone but various alleles are known to confer distinct patterns of pigmentation. R(Lc) is associated with unique pigmentation of leaves and R(Sn) with unique pigmentation of the scutellar node. One state of R is associated with pigmentation of the whole plant (R(P)), while another is associated with pigmentation of the seeds (R(S)).

Alleles of the unlinked B locus (in corn located on chromosome 2) rarely condition pigmentation of the aleurone, but are frequently associated with pigmentation of mature plant parts. The B-peru allele however, pigments the aleurone (like R(S)). Analysis at the molecular level has confirmed that the R and B loci are duplicate genes.

In order that the R and B loci can color a particular tissue, the appropriate allele of C1 or P1 loci also needs to be present. The C1 and C1-S alleles, for instance, pigment the aleurone when combined with the suitable R or B allele.

25

Alleles of the C1 locus have been cloned and sequenced. Of particular interest are C1 (Paz-Ares et al,

1987, EMBO J. 6:3553-3558) and C1-S (Schleffer et al, 1994, Mol.Gen.Genet. 242:40-48). Analysis of the sequences revealed the presence of two introns in the coding region of the gene. The protein encoded by the C1 and C1-S alleles shares homology with myb proto-oncogenes and is known to be a nuclear protein with DNA-binding capacity acting as transcriptional activators.

The cDNA of the B-peru allele has also been analyzed and sequenced (Radicella et al, 1991, Plant Mol. Biol. 17:127-130). Genomic sequences of B-peru were also isolated and characterized based on the homology between R and B (Chandler et al., 1989, the Plant Cell 1:1175-1183; Radicella et al., 1992, Genes & Development 6:2152-2164). The tissue specificity of anthocyanin production of two different B alleles was shown to be due to differences in the promoter and untranslated leader sequences (Radicella et al, 1992, supra).

Various alleles of the R gene family have also been characterized at the molecular level, e.g. Lc (Ludwig et al, 1989, PNAS 86:7092-7096), R-nj, responsible for pigmentation of the crown of the kernel (Dellaporta et al, 1988, In "Chromosome Structure and Function,: Impact of New Concepts, 18th Stadeler Genetics Symposium, Gustafson and Appels, eds. (New York, Plenum press, pp. 263-282)), Sn (Consonni et al, 1992, Nucl. Acids. Res.

20:373), and R(S) (Perrot and Cone, 1989, Nucl. Acids. Res. 17:8003).

The proteins encoded by the B and R genes share homology with myc proto-oncogenes and have characteristics of transcriptional activators.

It has been shown that various structural and regulatory genes introduced in maize tissues by microprojectiles operate in a manner similar to the endogenous loci and can complement genotypes which are deficient in the introduced genes (Klein et al., 1989, PNAS 86:6681-6685; Goff et al., 1990, EMBO J. 9:2517-2522). The Lc gene was also used as a visible marker for plant transformation (Ludwig et al., 1990, Science 247:449- 450). Apart from the above other genes involved in anthocyanin biosynthesis have been cloned (Cone, 1994, In 'The Maize Handbook', Freeling and Walbot eds., Springer Verlag New York Inc., p. 282-285).

In Barley, Falk et al (1981, In Barley Genetics IV, proceedings of the 4th International Barley Genetics symposium, Edinburgh University press, Edinburgh, pp. 778-785) have reported the coupling of a male-sterile gene to a xenia-expressing shrunken endosperm gene which makes it possible to select seeds, before planting, that will produce male-sterile plants. Problems associated with such proposal include complete linkage of the two genes

(Stoskopf, 1993, Plant Breeding : Theory and Practice, Westview Press, Boulder, San Francisco, Oxford). In sweetcorn, a genetic system to produce hybrid corn seeds without detassling, which utilizes the closely linked genes y (white endosperm) and ms (male sterility) was suggested but was never used because of contamination from 5% recombination. Galinat (1975, J.Hered. 66:387-388) described a two-step seed production scheme that resolved this problem by using electronic color sorters to separate yellow from white kernels . This approach has not been utilized commercially (Kankis and Davis, 1986, in « Breeding Vegetable Crops », the Avi Publishing Company Inc. Westport, Connecticut, U.S.A., p. 498).

EP 0,198,288 and US Patent 4,717,219 describe methods for linking marker genes (which can be visible markers or dominant conditional markers) to endogenous nuclear loci containing nuclear male-sterility genotypes.

EP 412,911 describes foreign restorer genes (e.g. barstar coding region under control of a stamen-specific promoter) that are linked to marker genes, including herbicide resistance genes and genes coding for pigments (e.g. the A1 gene) under control of a promoter which directs expression in specific cells, such as petal cells, leaf cells or seed cells, preferably in the outer layer of the seed.

Summary of the Invention

The invention concerns a maintainer plant consisting essentially of cells which comprise in their genome:

- 5 - a homozygous male-sterility genotype at a first genetic locus; and
- a color-linked restorer genotype at a second genetic locus, which is heterozygous (Rf/-) for a foreign DNA Rf comprising:
 - 10 a) a fertility-restorer gene capable of preventing the phenotypic expression of said male-sterility genotype, and
 - 15 b) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of said plant and which is capable of producing anthocyanin at least in the seeds of said plant, so that anthocyanin production in the seeds
 - 20 is visible externally.

The invention also concerns an anthocyanin regulatory gene which is a shortened R, B or C1 gene or a combination of shortened R, B or C1 genes which is

25 functional for conditioning and regulating anthocyanin production in the aleurone.

The invention also includes a DNA such as a plasmid comprising a fertility-restorer gene capable of preventing the phenotypic expression of a male-sterility genotype in a plant and at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of a plant and which is capable of producing anthocyanin at least in the seeds of a plant, so that anthocyanin production in the seeds is visible externally.

Also within the scope of the invention is a process to maintain a line of male-sterile plants, which comprises the following steps:

i) crossing:

a) a male-sterile parent plant of said line having, in a first genetic locus, a homozygous male-sterility genotype, and

b) a maintainer parent plant of said line consisting essentially of cells which comprise, stably integrated in their nuclear genome:

- a homozygous male-sterility genotype at a first genetic locus; and

- a colored-linked restorer genotype at a second genetic locus, which is heterozygous for a foreign DNA comprising:

i) a fertility-restorer gene capable of preventing the phenotypic expression of said male-sterility genotype, and

5

ii) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of said plant which is capable of producing anthocyanin at least in the seeds of said plant, so that anthocyanin production in the seeds is visible externally,

10

ii) obtaining the seeds from said parent plants, and

15

iii) separating on the basis of color, the seeds in which no anthocyanin is produced and which grow into male-sterile parent plants.

20

Preferably, the genome of the male-sterile parent plant does not contain at least one anthocyanin regulatory gene necessary for the regulation of anthocyanin biosynthesis in seeds of this plant to produce externally visible anthocyanin in the seeds. In one embodiment of the invention, the genome of the male-sterile parent plant contains a first anthocyanin regulatory gene and the

25

genome of the maintainer plant a second anthocyanin regulatory gene which, when present with the first anthocyanin regulatory gene in the genome of a plant, is capable of conditioning the production of externally visible anthocyanin in seeds.

The invention also concerns a process to maintain a line of maintainer plants, which comprises the following steps:

i) crossing:

a) a male-sterile parent plant as described previously, and

b) a maintainer parent plant as described previously,

ii) obtaining the seeds from said male-sterile parent plant, and

iii) separating on the basis of color, the seeds in which anthocyanin is produced and which grow into maintainer parent plants.

The invention also relates to a kit for maintaining a line of male-sterile or maintainer plants, said kit comprising:

- 5 a) a male-sterile parent plant of said line as described previously, having, in a first genetic locus, a homozygous male-sterility genotype and which is incapable of producing externally visible anthocyanin in seeds, and
- 10 b) a maintainer parent plant of said line as described previously.

Also within the scope of the invention is a process to
15 maintain the kit described previously which comprises:

- crossing said male-sterile parent plant with said maintainer parent plant;
- obtaining the seeds from said male-sterile parent
20 plants and optionally the seeds from said maintainer parent plant in which no anthocyanin is produced; and
- optionally growing said seeds into male-sterile parent plants and maintainer parent plants.

25 As mentioned above, the present invention provides means to maintain a line of male-sterile plants, particularly corn or wheat plants. These means can be in

the form of a process which comprises the following steps:

5 i) crossing A) a first parent plant of said line, which is male-sterile, and which is genetically characterized by the absence of at least one anthocyanin regulatory gene thereby being incapable of producing anthocyanin in seeds, particularly in the aleurone layer, and also by having at a first genetic locus a homozygous male-sterility genotype, and B) a second parent plant of said
10 line, which is male-fertile, and which is genetically characterized by having at said first genetic locus, said homozygous male- sterility genotype, and at a separate second genetic locus the genotype Rf/-,

15 whereby,

Rf is a foreign chimeric DNA (the "color-linked restorer gene") stably integrated in the nuclear genome
20 of said plant which comprises:

- a) a fertility-restorer gene that is capable of preventing the phenotypic expression, i.e. the male- sterility, of said male-sterility genotype.
- b) said at least one anthocyanin regulatory gene (the
25 "color gene") involved in the regulation of the anthocyanin biosynthesis in cells of seeds of said cereal plant which is capable of producing

anthocyanin at least in the seeds, particularly in the aleurone, of said cereal plant,

ii) obtaining the seeds from said first parent plants

iii) separating, on the basis of color, the seeds in

5 which no anthocyanin is produced and in which the genotype at said first genetic locus is said homozygous male-sterility genotype and the genotype at said second genetic locus is -/-, and the seeds in which anthocyanin is produced and in which the genotype at said first
10 genetic locus is said homozygous male-sterility genotype and the genotype at said second genetic locus is Rf/-.

Of particular interest in the invention is a second parent plant in which said at least one anthocyanin
15 regulatory gene comprises a gene derived from a genomic clone of an R or B gene, particularly an R or B gene that conditions anthocyanin production in the aleurone, preferably the B-peru allele (e.g. the shortened B-peru gene in pCOL13), and/or comprises a gene derived from a
20 genomic clone of the Cl gene (e.g. the gene with the sequence of SEQ ID NO 1 or SEQ ID NO 5) or the Cl-S gene.

The first genetic locus can be endogenous to plants of said line (in which case the homozygous male-sterility
25 genotype will be m/m), but is preferably a foreign locus with genotype S/S in which S is a foreign DNA which, when

expressed in a plant is capable of rendering the plant male-sterile. A preferred foreign DNA comprises at least:

s1) a male-sterility DNA encoding a RNA, protein or polypeptide which, when produced or overproduced in a cell of the plant, significantly disturbs the metabolism, functioning and/or development of the cell, and,

s2) a sterility promoter capable of directing expression of the male-sterility DNA selectively in stamen cells, preferably tapetum cells, of the plant; the male-sterility DNA being in the same transcriptional unit as, and under the control of, the sterility promoter.

In case such a foreign male-sterility genotype is used, the fertility-restorer gene in the foreign DNA Rf preferably comprises at least:

a1) a fertility-restorer DNA encoding a restorer RNA, protein or polypeptide which, when produced or overproduced in the same stamen cells as said male-sterility gene S, prevents the phenotypic expression of said foreign male-sterility genotype comprising S, and,

a2) a restorer promoter capable of directing expression of the fertility-restorer DNA at least in the same stamen cells in which said male-sterility gene S is expressed, so that the phenotypic expression of said male-sterility gene is prevented; the fertility-

restorer DNA being in the same transcriptional unit
as, and under the control of, the restorer promoter.
In case of an endogenous male-sterility genotype which is
homozygous for the recessive male-sterility allele m, the
fertility restorer gene is preferably a DNA comprising
the dominant allele M of said locus.

The present invention also provides the novel foreign
chimeric DNA Rf as used in the second parent plants,
plasmids comprising these chimeric genes, and host cells
comprising these plasmids.

The present invention also provides the shortened B-
peru gene in pCOL13 (SEQ ID NO 6) and the shortened C1
gene, particularly the EcoRI-SfiI fragment of pCOL9 of
SEQ ID NO 5.

The present invention further provides plants the
nuclear genome of which is transformed with the foreign
chimeric DNA Rf, particularly the second parent plant.

Detailed Description of the Invention

A male-sterile plant is a plant of a given plant
species which is male-sterile due to expression of a
male-sterility genotype such as a foreign male-sterility
genotype containing a male-sterility gene. A restorer
plant is a plant of the same plant species that contains

within its genome at least one fertility-restorer gene that is able to restore the male fertility in those offspring obtained from a cross between a male-sterile plant and a restorer plant and containing both a male-sterility genotype and a fertility-restorer gene. A restored plant is a plant of the same species that is male-fertile and that contains within its genome a male-sterility genotype and a fertility-restorer gene.

A line is the progeny of a given individual plant.

A gene as used herein is generally understood to comprise at least one coding region coding for an RNA, protein or polypeptide which is operably linked to suitable promoter and 3' regulatory sequences. A structural gene is a gene whose product is a e.g. an enzyme, a structural protein, tRNA or rRNA. For example anthocyanin structural genes encode enzymes (e.g. chalcone synthase) directly involved in the biosynthesis of anthocyanins in plant cells. A regulatory gene is a gene which encodes a regulator protein which regulates the transcription of one or more structural genes. For example the R, B, and Cl genes are regulatory genes that regulate transcription of anthocyanin structural genes.

For the purpose of this invention the expression of a gene, such as a chimeric gene, will mean that the promoter of the gene directs transcription of a DNA into a mRNA which is biologically active i.e. which is either capable of interacting with another RNA, or which is

capable of being translated into a biologically active polypeptide or protein.

The phenotype is the external appearance of the expression (or lack of expression) of a genotype i.e. of a gene or set of genes (e.g. male-sterility, seed color, presence of protein or RNA in specific plant tissues etc.)

As used herein, a genetic locus is the position of a given gene in the nuclear genome, i.e. in a particular chromosome, of a plant. Two loci can be on different chromosomes and will segregate independently. Two loci can be located on the same chromosome and are then generally considered as being linked (unless sufficient recombination can occur between them).

An endogenous locus is a locus which is naturally present in a plant. A foreign locus is a locus which is formed in the plant because of the introduction, by means of genetic transformation, of a foreign DNA.

In diploid plants, as in any other diploid organisms, two copies of a gene are present at any autosomal locus. Any gene can be present in the nuclear genome in several variant states designated as alleles. If two identical alleles are present at a locus that locus is designated as being homozygous, if different alleles are present, the locus is designated as being heterozygous. The allelic composition of a locus, or a set of loci, is the

genotype. Any allele at a locus is generally represented by a separate symbol (e.g. M and m, S and -, - representing the absence of the gene). A foreign locus is generally characterized by the presence and/or absence of a foreign DNA. A heterozygous genotype in which one allele corresponds to the absence of the foreign DNA is also designated as hemizygous (e.g. Rf/-). A dominant allele is generally represented by a capital letter and is usually associated with the presence of a biologically active gene product (e.g. a protein) and an observable phenotypic effect (e.g. R indicates the production of an active regulator protein and under appropriate conditions anthocyanin production in a given tissue while r indicates that no active regulator protein is produced possibly leading to absence of anthocyanin production).

A plant can be genetically characterized by identification of the allelic state of at least one genetic locus.

The genotype of any given locus can be designated by the symbols for the two alleles that are present at the locus (e.g. M/m or m/m or S/-). The genotype of two unlinked loci can be represented as a sequence of the genotype of each locus (e.g. S/S,Rf/-)

The nuclear male-sterility genotype as used in this invention refers to the genotype of at least one locus,

preferably only one locus, in the nuclear genome of a plant (the "male-sterility locus") the allelic composition of which may result in male sterility in the plant. A male-sterility locus may be endogenous to the plant, but it is generally preferred that it is foreign to the plant.

Foreign male-sterility loci are those in which the allele responsible for male sterility is a foreign DNA sequence S (the "male-sterility gene") which when expressed in cells of the plant make the plant male-sterile without otherwise substantially affecting the growth and development of the plant. Such male-sterility gene preferably comprises at least:

- s1) a male-sterility DNA encoding a sterility RNA, protein or polypeptide which, when produced or overproduced in a stamen cell of the plant, significantly disturbs the metabolism, functioning and/or development of the stamen cell, and,
- s2) a sterility promoter capable of directing expression of the male-sterility DNA selectively in stamen cells (e.g. anther cells or tapetum cells) of the plant; the male-sterility DNA being in the same transcriptional unit as, and under the control of, the sterility promoter.

The male-sterility locus preferably also comprises in the same genetic locus at least one first marker gene T which comprises at least:

- 5 t1) a first marker DNA encoding a first marker RNA, protein or polypeptide which, when present at least in a specific tissue or specific cells of the plant, renders the plant easily separable from other plants which do not contain the first marker RNA, protein or polypeptide encoded by the first marker DNA at least
10 in the specific tissue or specific cells, and,
- t2) a first marker promoter capable of directing expression of the first marker DNA at least in the specific tissue or specific cells: the first marker DNA being in the same transcriptional unit as, and
15 under the control of, the first marker promoter.

Such male-sterility gene is always a dominant allele at such a foreign male-sterility locus. The recessive allele corresponds to the absence of the male-sterility
20 gene in the nuclear genome of the plant.

Male-sterility DNAs and sterility promoters that can be used in the male-sterility genes in the first parent line of this invention have been described before (EP
25 0,344,029 and EP 0,412,911). For the purpose of this invention the expression of the male-sterility gene in a plant cell should be able to be inhibited or repressed

for instance by means of expression of a suitable fertility-restorer gene in the same plant cell. In this regard a particular useful male-sterility DNA codes for barnase (Hartley, J.Mol. Biol. 1988 202:913). The sterility promoter can be any promoter but it should at least be active in stamen cells, particularly tapetum cells. Particularly useful sterility promoters are promoters that are selectively active in stamen cells, such as the tapetum-specific promoters of the TA29 gene of Nicotiana tabacum (EP 0,344,029) which can be used in tobacco, oilseed rape, lettuce, cichory, corn, rice, wheat and other plant species; the PT72, the PT42 and PE1 promoters from rice which can be used in rice, corn, wheat, and other plant species (WO 92/13956) ; the PCA55 promoter from corn which can be used in corn, rice, wheat and other plant species (WO 92/13957); and the A9 promoter of a tapetum- specific gene of Arabidopsis thaliana (Wyatt et al., 1992, Plant Mol. Biol. 19:611-922). However, the sterility promoter may also direct expression of the sterility DNA in cells outside the stamen; particularly if the effect of expression of the male-sterility DNA is such that it will specifically disturb the metabolism, functioning and/or development of stamen cells so that no viable pollen is produced. One example of such a male-sterility DNA is the DNA coding for an antisense RNA which is complementary to the mRNA of the chalcone synthase gene (van der Meer et al (1992)

The Plant Cell 4:253-262). In this respect a useful promoter is the 35S promoter (see EP 0,344,029), particularly a 35S promoter that is modified to have enhanced activity in tapetum cells as described by van der Meer et al (1992) The Plant Cell 4:253-262 (the "35S-tap promoter").

A preferred endogenous male-sterility locus is one in which a recessive allele (hereinafter designated as m) in homozygous condition (m/m) results in male sterility. At such loci male fertility is encoded by a corresponding dominant allele (M). In many plant species such endogenous male- sterility loci are known (see Kaul, 1988, supra (in corn see also recent issues of Maize Genetics Cooperation Newsletter, published by Department of Agronomy and U.S. Department of Agriculture, University Of Missouri, Columbia, Missouri, U.S.A.)). The DNA sequences in the nuclear genome of the plant corresponding to m and M alleles can be identified by gene tagging i.e. by insertional mutagenesis using transposons, or by means of T-DNA integration (see e.g. Wienand and Saedler, 1987, In 'Plant DNA Infectious Agents', Ed. by T.H.Hohn and J.Schell, Springer Verlag Wien New York, p. 205; Shepherd, 1988, In 'Plant Molecular Biology: a Practical Approach', IRL Press, p. 187; Teeri et al., 1986, EMBO J. 5:1755). It will be evident that in the first and second parent plant of this

invention S/S can be replaced by m/m without affecting the outcome of the process. Indeed, one feature of the process of this invention is that the male-sterility locus is homozygous thus allowing the use of 'recessive' male-sterility alleles.

Fertility-restorer DNAs that can be used in the fertility restorer gene in the second parent line of this invention have been described before (EP 0,412,911).

In this regard, fertility-restorer genes in which the fertility-restorer DNA encodes barstar (Hartley, J.Mol. Biol. 1988 202:913) are particularly useful to inhibit the expression of a male-sterility DNA that encodes barnase. In this regard it is believed that a fertility-restorer DNA that codes for a mutant of the barstar protein, i.e. one in which the Cysteine residue at position 40 in the protein is replaced by serine (Hartley, 1989, TIBS 14:450), functions better in restoring the fertility in the restored plants of some species.

In principle any promoter can be used as a restorer promoter in the fertility restorer gene in the second parent line of this invention. The only prerequisite is that such second parent plant, which contains both the color gene and the fertility-restorer gene, should be phenotypically normal and male-fertile. This requires that the restorer promoter in the fertility-restorer gene

should be at least active in those cells of a plant of the same species in which the sterility promoter of the corresponding male-sterility gene can direct expression of the male-sterility DNA. In this regard it will be preferred that the sterility promoter and the restorer promoter are the same; they can for example be both stamen-specific promoters (e.g. the TA29 promoter or the CA55 promoter) or they can be both constitutive promoters (such as the 35S or 35S-tap promoter). However, the sterility promoter may be active only in stamen cells while the restorer promoter is also active in other cells. For instance, the sterility promoter can be a stamen-specific (such as the TA29 or CA55 promoter) while the restorer promoter is the 35S-tap promoter.

When the male sterility to be restored is due to the male- sterility genotype at an endogenous male-sterility locus being homozygous for a recessive allele *m*, it is preferred that the fertility-restorer gene is the dominant allele of that male- sterility locus, preferably under control of its own promoter. The DNA corresponding to such a dominant allele, including its natural promoter can be isolated from the nuclear genome of the plant by means of gene tagging as described above.

The nature of the color gene that is used in the color- linked restorer gene in the second parent plant of

this invention depends upon the genotype of the untransformed plants of the same line. Preferably, only cereal plants with a genotype that does not condition externally visible anthocyanin production in seeds, particularly in the aleurone can be used to produce the second parent plants. These plants usually have a genotype in which no functional copy of a suitable regulatory gene such as the R or B gene, and/or the C1 gene, is present.

In corn, for instance, all of the currently used inbred lines in the U.S.A. are r-r (pink anthers, leaf tips, plant base) or r-g (green) and most of these are c1 and pl; at the B- locus the B-peru allele is very rare (Coe et al, 1988, In 'Corn and Corn Improvement', 3rd edition, G.F.Sprague and J.W. Dudley, eds. America Science of Agronomy, Inc. Publishers, Madison, Wisconsin, U.S.A.). The result is that no anthocyanins are produced in the aleurone of these lines and that the kernels are yellow. This requires that when these lines are transformed with a color-linked restorer gene, the color gene should consist of a functional R or B gene which conditions anthocyanin production in aleurone, and usually also a functional C1 gene capable of conditioning anthocyanin production in aleurone.

A useful R or B gene is the B-peru gene, but of course also other R genes could be used such as the R(S) gene (Perrot and Cone, 1989, Nucl. Acids Res. 17:8003).

In this regard a gene derived from genomic clones of the B-peru gene (Chandler et al, 1989, The Plant Cell 1:1175-1183) is believed to be particularly useful. However the length of this genomic DNA (11 kbp) renders its practical manipulation and use for transformation by direct gene transfer, difficult, certainly in combination with other genes such as the restorer gene and the C1 gene.

In one inventive aspect of this invention it was found that the B-peru gene could be considerably shortened while still retaining, under appropriate conditions, its capability of conditioning anthocyanin production in the aleurone of seeds of cereal plants such as corn. A preferred shortened B-peru gene is that of Example 2.2 and which is contained in plasmid pCOL13 (deposited under accession number LMBP 3041).

A useful C1 gene is the genomic clone as described by Paz-Ares et al, 1987, EMBO J. 6:3553-3558. However the length of this genomic DNA (4 kbp) precludes its practical manipulation and use for transformation by direct gene transfer, certainly in combination with other genes such as the restorer gene and the B-peru gene. Nevertheless other variants of the C1 gene can also be used. In this regard Scheffler et al, 1994, Mol.Gen.Genet. 242:40-48 have described the C1-S allele which differs from the C1 allele of Paz-Ares et al, supra by a few nucleotides in the promoter region near the CAAT

box and which is dominant to the wild-type allele (C1) and shows enhanced pigmentation. The C1-S gene can be easily used in this invention by appropriate changes in the C1 gene. For example the TGCAG at positions 935 to 939 in SEQ ID NO 1 (respectively at positions 884-888 in SEQ ID NO 5) can be easily changed to TTAGG yielding a C1-S allele (respectively pCOL9S).

In one inventive aspect of this invention it was found that the C1 gene (and the C1-S gene) could be considerably shortened while still retaining, under appropriate conditions, its capability of conditioning anthocyanin production in the aleurone of seeds of cereal plants such as corn. Preferred shortened C1 genes for instance are those of Example 2.1 such as comprised in pCOL9 which has the sequence of SEQ ID NO 5, particularly as comprised between the EcoRI and SfiI sites of pCOL9, and the corresponding shortened C1-S gene in pCOL9S.

The transcribed region of the shortened B-peru and C1 genes still contain some small introns which can also be deleted without affecting the function of the genes. It is also believed that the shortened B-peru and C1 genes can be somewhat further truncated at their 5' and 3' ends, without affecting their expression in aleurone. In particular it is believed that the sequence between positions 1 and 3272 of SEQ ID NO 6 can also be used as a suitable B-peru gene. It is also believed that this gene

can still be truncated at its 3' end down to a position between nucleotides 2940 and 3000 of SEQ ID No. 6.

Although the use of genomic sequences of the B-peru gene and the C1 gene, particularly the shortened B-peru and/or the shortened C1 of C1-S genes, is preferred, chimeric R, B, or C1 genes can also be used. For instance a chimeric gene can be used which comprises the coding region (e.g. obtained from the cDNA) of any functional R or B gene (i.e. which conditions anthocyanin production anywhere in the plant) which is operably linked to the promoter region of a R or B gene which conditions anthocyanin production in the aleurone (such as R(S) or B-peru). Since the presence of anthocyanin does not negatively affect growth, development and functioning of plant cells, a constitutive promoter (e.g. the 35S promoter), or a promoter which directs expression at least in the aleurone can also be used in such a chimeric gene. In this regard the promoter of the C1 gene can also be used to direct expression of a DNA comprising the coding region of suitable R or B gene, particularly the B-peru gene.

Similarly the coding region (e.g. obtained from cDNA) of the C1 gene can be operably linked to the promoter of a gene that directs expression at least in the aleurone. In this regard, the promoter of the B-peru gene can also be used to direct expression of a DNA comprising the

coding region of a suitable C1 gene such as that of the C1 gene of SEQ ID No. 1 or of the C1-S gene.

5 In another inventive aspect of the invention it was found that the the promoters comprised in DNAs characterized by the sequences between positions 1 to 1077, particularly between positions 447 and 1077, quite particularly between positions 447 and 1061 of SEQ ID NO 1, between positions 396 and 1026 of SEQ ID NO 5, and between positions 1 to 575, particularly between position 10 1 to 188 of SEQ ID NO 6 are promoters that predominantly, if not selectively, direct expression of any DNA, preferably a heterologous DNA in the aleurone layer of the seeds of plants.

15 Of course in those lines in which a functional C1 gene is already present in the genome the color gene can consist only of a suitable functional R or B gene (or a chimeric alternative). Alternatively if a line contains already a functional R or B gene which can condition 20 anthocyanin production in the aleurone, but no functional C1 gene, only a functional C1 gene is required as a color gene.

25 It is believed that the color genes of this invention are especially useful in cereal plants, and that they are of particular use in corn and wheat, and certainly in corn.

For the purposes of this invention it is preferred that, in the second parent plants the "Rf" locus and the male- sterility (e.g. "S") locus are not linked and segregate separately.

In the second parent plant, the fertility restorer gene, the B-peru gene and the Cl gene are preferably closely linked. This can of course be achieved by introducing these genes in the nuclear genome of the plants as a single transforming foreign DNA (the Rf DNA) thus forming a foreign Rf locus. Alternatively, the fertility restorer gene and the color gene can be separately introduced by cotransformation which usually results in single locus insertions in the plant genome.

The color-linked restorer gene Rf as used in the second parent plant preferably also comprises at least c) a second marker gene which comprises at least:

c1) a second marker DNA encoding a second marker RNA, protein or polypeptide which, when present at least in a specific tissue or specific cells of the plant, renders the plant easily separable from other plants which do not contain the second marker RNA, protein or polypeptide encoded by the second marker DNA at least in the specific tissue or specific cells, and,

c2) a second marker promoter capable of directing expression of the second marker DNA at least in the specific tissue or specific cells: the second marker DNA being in the same transcriptional unit as, and under the control of, the second marker promoter.

First and second marker DNAs and first and second marker promoters that can be used in the first and second marker genes of this invention are also well known (EP 0,344,029; EP 0,412,911). In this regard it is preferred that the first and second marker DNA are different, although the first and second marker promoter may be the same.

Foreign DNA such as the fertility-restorer gene, the foreign male-sterility gene, the B-peru and the C1 genes, or the first or second marker gene preferably also are provided with suitable 3' transcription regulation sequences and polyadenylation signals, downstream (i.e. 3') from their coding sequence i.e. respectively the fertility-restorer DNA, the male-sterility DNA, the coding region of a color gene (such as a B-peru gene and/or a C1 gene) or the first or second marker DNA. In this regard either foreign or endogenous transcription 3' end formation and polyadenylation signals suitable for obtaining expression of the chimeric gene can be used. For example, the foreign 3' untranslated ends of genes,

such as gene 7 (Velten and Schell (1985) Nucl. Acids Res. 13:6998), the octopine synthase gene (De Greve et al., 1982, J.Mol. Appl. Genet. 1:499; Gielen et al (1983) EMBO J. 3:835; Ingelbrecht et al., 1989, The Plant Cell 1:671) and the nopaline synthase gene of the T-DNA region of Agrobacterium tumefaciens Ti-plasmid (De Picker et al., 1982, J.Mol. Appl. Genet. 1:561), or the chalcon synthase gene (Sommer and Saedler, 1986, Mol.Gen.Genet. 202:429-434), or the CaMV 19S/35S transcription unit (Mogen et al., 1990, The Plant Cell 2:1261-1272) can be used. However, it is preferred that the color genes in this invention carry their endogenous transcription 3' end formation and polyadenylation signals.

The fertility-restorer gene, the male-sterility gene, the color gene or the first or second marker gene in accordance with the present invention are generally foreign DNAs, preferably foreign chimeric DNA. In this regard "foreign" and "chimeric" with regard to such DNAs have the same meanings as described in EP 0,344,029 and EP 0,412,911.

The cell of a plant, particularly a plant capable of being infected with Agrobacterium such as most dicotyledonous plants (e.g. Brassica napus) and some monocotyledonous plants, can be transformed using a vector that is a disarmed Ti-plasmid containing the male-

sterility gene, the color linked restorer gene or both and carried by Agrobacterium. This transformation can be carried out using the procedures described, for example, in EP 0,116,718 and EP 0,270,822. Preferred Ti-plasmid vectors contain the foreign DNA between the border sequences, or at least located to the left of the right border sequence, of the T-DNA of the Ti-plasmid. Of course, other types of vectors can be used to transform the plant cell, using procedures such as direct gene transfer (as described, for example, in EP 0,233,247), pollen mediated transformation (as described, for example, in EP 0,270,356, PCT patent publication "WO" 85/01856, and US patent 4,684,611), plant RNA virus-mediated transformation (as described, for example, in EP 0,067,553 and US patent 4,407,956) and liposome-mediated transformation (as described, for example, in US patent 4,536,475). Cells of monocotyledonous plants such as the major cereals including corn, rice, wheat, barley, and rye, can be transformed (e.g. by electroporation) using wounded or enzyme-degraded intact tissues capable of forming compact embryogenic callus (such as immature embryos in corn), or the embryogenic callus (such as type I callus in corn) obtained thereof, as described in WO 92/09696. In case the plant to be transformed is corn, other recently developed methods can also be used such as, for example, the method described for certain lines of corn by Fromm et al., 1990, Bio/Technology 8:833;

Gordon-Kamm et al., 1990, Bio/Technology 2:603 and Gould et al., 1991, Plant Physiol. 95:426. In case the plant to be transformed is rice, recently developed methods can also be used such as, for example, the method described for certain lines of rice by Shimamoto et al., 1989, Nature 338:274; Datta et al., 1990, Bio/Technology 8:736; and Hayashimoto et al., 1990, Plant Physiol. 93:857.

The transformed cell can be regenerated into a mature plant and the resulting transformed plant can be used in a conventional breeding scheme to produce more transformed plants with the same characteristics or to introduce the male- sterility gene, the color-linked restorer gene (or both), in other varieties of the same related plant species. Seeds obtained from the transformed plants contain the chimeric gene(s) of this invention as a stable genomic insert. Thus the male- sterility gene, or the color-linked restorer gene of this invention when introduced into a particular line of a plant species can always be introduced into any other line by backcrossing.

The first parent plant of this invention contains the male-sterility gene as a stable insert in its nuclear genome (i.e. it is a male-sterile plant). For the purposes of this invention it is preferred that the first parent plant contains the male-sterility gene in

homozygous condition so that it transmits the gene to all of its progeny.

The second parent plant of this invention contains the male-sterility gene and the color-linked restorer gene as stable inserts in its nuclear genome (i.e. it is a restored plant). It is preferred that the male-sterility gene be in homozygous condition so that the second parent plant transmits the gene to all of its progeny and that the color-linked restorer gene be in heterozygous condition so that the second parent plant transmits the gene to only half of its progeny.

It is preferred that the first and second parent plants are produced from the same untransformed line of a plant species, particularly from the same inbred line of that species.

The first and second parent plants of this invention have the particular advantage that seeds of such plants can be maintained indefinitely, and can be amplified to any desired amount (e.g. by continuous crossing of the two plant lines).

The color genes of this invention can be used as marker gene in any situation in which it is worthwhile to detect the presence of a foreign DNA (i.e. a transgene) in seeds of a transformed plant in order to isolate seeds which possess the foreign DNA. In this regard virtually

any foreign DNA, particularly a chimeric gene can be linked to the color gene.

5 Examples of such foreign DNAs are genes coding for insecticidal (e.g. from Bacillus thuringiensis), fungicidal or nematocidal proteins. Similarly the color-gene can be linked to a foreign DNA which is the male-sterility gene as used in this invention.

10 However, the color genes are believed to be of particular use in the process of this invention in which they are present in a foreign DNA which comprises a fertility restorer gene (such as the barstar gene of Bacillus amyloliquefaciens) under control of a stamen-specific promoter (such as PTA29). In appropriate
15 conditions the use of the color genes allows the easy separation of harvested seeds that will grow into male-sterile plants, and harvested seeds that will grow into male-fertile plants. In this regard the seeds are preferably harvested from male-sterile plants (the first
20 parent plants) that are homozygous at a male-sterility locus (such as a locus comprising the barnase gene under control of PTA29) and which have been pollinated by restorer plants (the second parent plants of this invention) which contain in their genome two unlinked
25 gene loci one of which comprises the same male-sterility locus which is homozygous for the same male-sterility gene while the other is a foreign locus which comprises

an appropriate fertility restorer gene (i.e. whose expression will counteract the expression of the male-sterility gene) and also the color gene of this invention, particularly an R or B gene that is expressed in the aleurone and/or a C1 gene, preferably the B-peru and C1 gene (e.g. as described in the examples). First and second parent plants can be essentially produced as described in the examples and as summarized in Figure 1. In step 8 of Figure 1 it is demonstrated that the crossing of the first and second parent plants of this invention will give rise in the progeny to about 50% new first parent (i.e. male-sterile) plants and about 50% new second parent (i.e. male-fertile) plants and that these two types of plants can already be separated at the seed stage on the basis of color. Red kernels will grow into male-fertile plants while yellow kernels will grow into male-sterile plants.

Thus a line of male-sterile first parent plants of this invention can be easily maintained by continued crossing with the second parent plants of this invention with, in each generation, harvesting the seeds from the male-sterile plants and separation of the yellow and red kernels. Of course in this way any desired amount of seed for foundation seed production of a particular line, such as an inbred line, can also be easily obtained.

The red and yellow seeds harvested from a cereal plant (e.g. the first parent plant of this invention) can

be separated manually. However, such separation can also be effected mechanically. A color sorting machine for corn kernel and other granular products is for instance available from Xeltron U.S. (Redmond, Washington, U.S.A.)

5

Unless otherwise indicated all experimental procedures for manipulating recombinant DNA were carried out by the standardized procedures described in Sambrook et al., 1989, "Molecular Cloning: a Laboratory Manual", Cold Spring Harbor Laboratory, and Ausubel et al, 1994, "Current Protocols in Molecular Biology", John Wiley & Sons.

10

The polymerase chain reactions ("PCR") were used to clone and/or amplify DNA fragments. PCR with overlap extension was used in order to construct chimeric genes (Horton et al, 1989, Gene 77:61-68; Ho et al, 1989, Gene 77:51-59).

15

All PCR reactions were performed under conventional conditions using the VentTM polymerase (Cat. No. 254L - Biolabs New England, Beverley, MA 01915, U.S.A.) isolated from Thermococcus litoralis (Neuner et al., 1990, Arch.Microbiol. 153:205-207). Oligonucleotides were designed according to known rules as outlined for example by Kramer and Fritz (1968, Methods in Enzymology 154:350), and synthesized by the phosphoramidite method (Beaucage and Caruthers, 1981, Tetrahedron Letters

20
25

22:1859) on an applied Biosystems 380A DNA synthesizer (Applied Biosystems B.V., Maarssen, Netherlands).

In the following examples, reference will be made to the following sequence listing and figures:

Sequence Listing

SEQ ID NO 1 : sequence of C1 gene

10 SEQ ID NO 2 : plasmid pTS256

SEQ ID NO 3 : EcoRI-HindIII region of pTS200 comprising the chimeric gene PCA55-barstar-3'nos (the omitted region of pTS200 is derived from pUC19.

15 SEQ ID NO 4 : oligonucleotide 1

SEQ ID NO 5 : pCOL9 containing the shortened C1 gene as a EcoRI-SfiI fragment

20 SEQ ID NO 6 : presumed sequence of the EcoRI-HindIII region of pCOL13 containing the shortened B-peru gene (the rest of the plasmid is pUC19). The stretch of N nucleotides corresponds to a region of approximate length which is derived from the genomic clone of the B-peru gene but for

25

which the sequence needs to be confirmed.

SEQ ID NO 7 : actual sequence of the EcoRI-HindIII region of pCOL13 containing the shortened B-peru gene (the rest of the plasmid is pUC19).

Figures

Figure 1 : Breeding scheme to obtain the first and second parent plants of this invention

Figure 2 : Schematic structure of pCOL25, pCOL26, pCOL27, pCOL28, pCOL100 and pDE110.

Examples

Example 1 : Construction of plasmids containing the male-sterility gene comprising the TA29 promoter and the barnase coding region

Plasmids useful for transformation of corn plants and carrying a male-sterility gene and a selectable marker gene have been described in WO 92/09696 and WO 92/00275.

Plasmid pVE107 contains the following chimeric genes:

1) PTA29-barnase-3'nos, i.e. a DNA coding for barnase of Bacillus amyloliquefaciens (barnase) operably linked to the stamen-specific promoter of the TA29 gene of

Nicotiana tabacum (PTA29) and the 3' regulatory sequence containing the polyadenylation signal of the nopaline synthase gene of Agrobacterium tumefaciens (3'nos), and 2) P35S-neo-3'ocs, i.e. the coding region of the gene of Tn5 of E.coli coding for neomycin phosphotransferase (neo) operably linked to the 35S promoter of Cauliflower Mosaic Virus (P35S) and the 3' regulatory sequence containing the polyadenylation signal of the octopine synthase gene of Agrobacterium tumefaciens (3'ocs).

Plasmid pVE108 contains the following chimeric genes: 1) PTA29-barnase-3'nos, and 2) P35S-bar-3'nos, i.e. the gene of Streptomyces hygroscopicus (EP 242236) coding for phosphinothricin acetyl transferase (bar) operably linked to the P35S and 3'nos.

PTA29-barnase-3'nos is an example of a foreign chimeric male-sterility gene (S) used in this invention.

Example 2 : Construction of a plasmid containing the color-linked restorer gene

2.1. Obtaining a shortened functional C1 gene

The C1 gene of maize was cloned from transposable-induced mutants and its sequence was reported (Paz-Ares, 1987, EMBO J. 6:3553-3558). This sequence is reproduced in SEQ ID NO. 1. Plasmid p36 (alternatively designated as pC1LC5kb and further designated as plasmid pXX036) comprising a C1 genomic clone was obtained from Dr. H. Saedler and Dr. U. Wienand of the Max-Planck Institut

für Züchtungsforschung, Köln, Germany. pXX036 was digested with SnaBI and HindIII, filled-in with Klenow, and selfligated, yielding plasmid pCOL9. pCOL9 corresponds to pUC19 (Yanisch-Perron et al, 1985, Gene 33:103-119) which contains, between its EcoRI and modified HindIII sites, the 2189 bp EcoRI-SnaBI fragment (corresponding to the sequence between positions 448 and 2637 of SEQ ID NO 1) of pXX036.

pXX036 was also digested with SfiI and HindIII and treated with Klenow to make blunt ends. After ligation the plasmid in which the DNA downstream from the SfiI site was deleted was designated as pCOL12.

The sequence TGCAG in pCOL9, corresponding to the sequence at positions 884 to 888 in SEQ ID NO 5, is changed to TTAGG, yielding pCOL9S which instead of a shortened C1 gene contains a shortened overexpressing C1-S gene (Schleffer et al, 1994, Mol.Gen.Genet. 242:40-48). A similar change is introduced in pCOL12, yielding pCOL12S.

2.2. Obtaining a shortened functional B-peru gene

Plasmid pBP2 (further designated as pXX004) is plasmid pTZ18U (Mead et al., 1986, Protein Engineering 1:67; U.S.Biochemical Corp.) containing the genomic clone of the B- peru gene. Plasmid p35SBPcDNA (further

designated as pXX002) is plasmid pMF6 (Goff et al, 1990, EMBO J. 9:2517-2522) containing the cDNA corresponding to the B-peru gene. Both plasmids were obtained from Dr. V. Chandler of the University of Oregon, Oregon, U.S.A. A
5 2660 bp sequence of the genomic clone around the translation initiation codon was reported (EMBL/Genbank/DDBJ databases; locus name ZMBPERUA, Accession number X70791; see also Radicella et al, 1992, Genes & Development 6:2152-2164). The sequence of the B-
10 peru cDNA was also reported (Radicella et al, 1991, Plant Mol. Biol. 17:127- 130).

Substantial amounts of 5' and 3' flanking sequences were deleted from pXX004, and the MluI-MunI fragment in the coding region of the genomic clone was replaced by
15 the 1615 bp MluI- MunI fragment of the cDNA clone. The resulting plasmid was designated as pCOL13 which was deposited at the Belgian Coordinated Collection of Microorganisms - LMBP Collection, Laboratory Molecular Biology, University of Ghent, K.L. Ledeganckstraat 35, B-
20 9000 Ghent, Belgium and was given the Accession Number LMBP 3041. A shortened but functional B-peru gene is contained in pCOL13 as an EcoRI-SalI fragment with an approximate length of 4 kbp (see SEQ ID NO 6).

25 2.3. Combining the C1 and B-peru genes

The C1 gene in pCOL9 and the B-peru gene in pCOL13 were then combined as follows. The 4 kbp EcoRI-SalI

fragment of pCOL13 was introduced between the EcoRI and SalI sites of the vector pBluescript II SK(-) (Stratagene), yielding #7 B SK(-). pCOL9 was digested with SfiI, treated with Klenow to fill in protruding ends, and further digested with EcoRI. The 1978 bp SfiI(Klenow)/EcoRI was then introduced between the EcoRI and SmaI sites of #7 B SK(-), yielding #7 B+C SK(-). Finally the XhoI site in the C1 sequence was removed as follows. The 950 bp EcoRI-SacII fragment of #7 B SK(-) (EcoRI site corresponding to the EcoRI site at position 1506 in SEQ ID NO 1; the SacII site from the pBluescript linker) was introduced between the EcoRI and SacII sites of the Phagescript Vector (Stratagene) to yield pCOL21. Single strands of pCOL21 were prepared and hybridized to the following synthetic oligonucleotide 1 (SEQ ID No. 4):

5'-CGT TTC TCG AAT CCG ACG AGG-3'

resulting in a silent change (CTCGAG -> CTCGAA) and removal of the XhoI site.

The 710 AatII-SacII fragment of #7 B SK(-) was then exchanged for the corresponding AatII-SacII fragment of the mutated pCOL21, yielding pCOL23.

pCOL23 was then linearized with SacII, treated with Klenow, and ligated to XhoI linker sequence (Stratagene), yielding pCOL24.

25

Using the same procedure as described above, the shortened C1-S gene of pCOL9S is combined with the

shortened B-peru gene of pCOL23, yielding plasmid pCOL24S.

2.4. Construction of vectors comprising the Cl and B-peru genes as well as male-sterility gene and a selectable marker gene

PTS256 is derived from pUC19 and contains the following two chimeric genes :1) P35S-bar-3'nos, and 2) PTA29-barstar- 3'nos, i.e. a DNA coding for barstar of Bacillus amyloliquefaciens (barstar or bar*) operably linked to PTA29 and 3'nos. The complete sequence of PTS256 is given in SEQ ID NO 2.

PTS200 is derived from pUC19 and contains the following two chimeric genes : 1) P35S-bar-3'nos, and 2) PCA55-barstar- 3'nos, i.e. barstar operably linked to the stamen-specific promoter PCA55 of Zea mays and 3'nos. The complete sequence of PTS200 is given in SEQ ID NO 3.

PTS256 was modified by the inclusion of NotI linkers (Stratagene) in both the unique SspI and SmaI sites, yielding PTS256NN. The shorter BspEI-SacII fragment of PTS256NN was then replaced by the shorter BspEI-SacII fragment of PTS200, yielding PTS256+200.

PTS256NN contains P35S-bar3'-nos and pTA29-barstar3'nos on a NotI cassette. PTS256NN+200 contains P35S-bar3'-nos and PCA55- barstar3'nos on a NotI cassette.

The NotI cassette of PTS256NN was introduced in the NotI site of pCOL24, yielding pCOL25 and pCOL26 which

differ with respect to the orientation of the P35S-bar3'-nos gene with respect to the shortened C1 gene (Figure 2).

5 The NotI cassette of pTS256NN+200 was introduced in the NotI site of pCOL24, yielding pCOL27 and pCOL28 which differ with respect to the orientation of the P35S-bar3'-nos gene with respect to the shortened C1 gene (Figure 2).

10 Plasmids pCOL25, pCOL26, pCOL27 or pCOL28 contain a color- linked restorer gene Rf and a selectable marker gene (P35S-bar- 3'nos). Rf comprises the shortened C1 and B-peru genes and a chimeric barstar gene (either PTA29-barstar-3'nos or PCA55- barstar-3'nos).

15 Plasmids pCOL25S, pCOL26S, pCOL27S or pCOL28S, containing the shortened C1-S gene instead of the shortened C1 gene, are obtained in a similar way using pCOL24S instead of pCOL24.

20 2.5. Construction of vectors comprising the C1 and B-peru genes as well as male-sterility gene

Plasmid pTS59 can be obtained from plasmid pTS256 (of SEQ ID NO 2) by replacing the fragment extending from positions 1 to 1470 (comprising the chimeric gene P35S-bar-3'nos) with the sequence TATGATA. Then NotI linkers (Stratagene) were introduced in the EcoRV and SmaI sites of pTS59; yielding pTS59NN. Finally the NotI fragment

25

comprising the chimeric gene PTA29-barstar-3'nos was introduced in the NotI site of #7 B+C SK(-), yielding pCOL100 (the general structure of pCOL100 and pDE110 is also presented in Figure 2).

5

2.6. Expression of shortened C1 and B-peru in aleurone in corn seeds

Dry seeds were incubated overnight in water at room temperature and were then peeled and sliced in half. Four
10 to six half kernels were placed with the cut side on wet filter paper and were bombarded with tungsten particles (diameter 0.7 μ m) which were coated with DNA.

Particle bombardment was essentially carried out using the particle gun and procedures as described by Zumbrunn
15 et al, 1989, Technique, 1:204-216. The tissue was placed at 10 cm from the stopping plate while a 100 μ m mesh was placed at 5 cm from the stopping plate.

- DNA of the following plasmids was used :
- 20 - pXX002 : B-peru cDNA under control of the 35S promoter
- pXX201 : C1 cDNA under control of the 35S promoter
- pCOL13 : shortened B-peru gene as described in Example 2.2
- pCOL12 : shortened C1 gene as described in Example 2.1
- pCOL100 : shortened B-peru and shortened C1 and PTA29-
25 barstar-3'nos as described in Example 2.5.

After bombardment the tissue was incubated for 2 days on wet filter paper at 27°C and was then checked for the presence of red spots indicating anthocyanin production.

Table 1

			pXX00	PXX201	pXX002 pXX201	pCOL13	pCOL12	pCOL100
H99	r	cl	-	-	+	nt	nt	nt
Pa91	r	cl	-	-	+	nt	nt	nt
B73	r	cl	-	-	+	nt	nt	+
inbred1	r	cl	-	-	+	nt	nt	nt
inbred2	r	cl	-	-	+	nt	nt	nt
inbred3	r	cl	-	-	+	nt	nt	nt
inbred4	r	cl	+	-	+	+	-	nt
inbred5	r	cl	+	-	+	+	-	nt
inbred6	r	cl	-	-	+	nt	nt	nt
inbred7	r	cl	-	-	+	nt	nt	nt
inbred8	r	cl	-	-	+	nt	nt	nt
inbred9	r	cl	+	-	+	+	-	nt
c-ruq	R	cl	-	+	+	-	+	nt

Note : + indicates that anthocyanin production was observed in at least one experiment; - indicates that no anthocyanin production was observed, nt = not tested.

The results for three public lines (H99, Pa91, B73) and 9 different, commercially important, proprietary inbred lines from various sources are shown in Table 1. The line c-ruq is a tester line which is homozygous for a C1 allele that is inactivated by insertion of a receptor for the regulator Uq (Cormack et al., 1988, Crop Sci. 28:941-944).

All lines which were r and c1 produced anthocyanin in the aleurone after introduction with both a functional B-peru and C1 gene. Lines which were R and c1 produced anthocyanin upon introduction of a functional C1 gene. Lines which were r and C1 produced anthocyanin upon introduction of a functional B-peru gene. This proves that the B-peru and C1 gene are sufficient for anthocyanin production in most corn lines. From the data in Table 1 it is also evident that even the shortened B-peru and C1 genes are still functional and are capable of producing anthocyanin in aleurone of corn lines with suitable genotypes.

Example 3 : Production of first parent corn plants by transformation of corn with the plasmids of example 1.

Corn plants of line H99, transformed with a male-sterility gene comprising a DNA encoding barnase of Bacillus amyloliquefaciens under control of the promoter of the TA29 gene of Nicotiana tabacum have been described in WO 92/09696. The transformed plants were shown to be male-sterile.

Example 4 : Production of second parent corn plants by transformation of corn with the plasmids of examples 2.

Corn inbred lines H99 and Pa91 are transformed using the procedures as described in WO 92/09696 but using plasmids pCOL25, pCOL26, pCOL27 or pCOL28 of Example 2.

Regenerated plants are selected that are male fertile and in which the shortened C1, the shortened B-peru gene, the P35S-bar-3'nos gene, and the PTA29- barstar-3'nos (or PCA55-barstar-3'nos) are expressed.

5 Alternatively the male-sterile plants of Example 3 (already containing the S gene) can be transformed with plasmids pCOL25, pCOL26, pCOL27 or pCOL28 of Example 2 on the condition that the S and Rf genes are linked to different selectable marker genes.

10

Similarly, transformed corn plants are obtained using plasmids pCOL25S, pCOL26S, pCOL27S or pCOL28S of Example 2.

15

In an alternative set of experiments the second parent plants of this invention were obtained by transforming corn plants of line H99, Pa91, and (Pa91xH99)x H99 with two separate plasmids one of which contained the color linked restorer gene (pCOL100), while the other contains an appropriate selectable marker gene such as a chimeric bar gene (pDE110) (alternatively a chimeric neo gene may also be used). pDE110 was described in WO 92/09696 and the construction of pCOL100 was described in Example 2.5.

20

25 In yet another set of experiments the second parent plants of this invention are obtained by transforming

corn plants with a purified fragment of the plasmids of example 2.4. Such purified fragment is obtained by digestion of the plasmids of example 2.4 with XhoI and subsequent purification using conventional procedures such as gel filtration.

Untransformed corn plants of lines H99 or Pa91 are detasseled and pollinated with pollen of the plants transformed with the Rf DNA. It is observed that the f gene segregates in a Mendelian way and that the seed that is harvested from these plants is colored and non-colored (yellow) in a 1:1 ratio. The red color of the seeds is correlated with the presence of the Rf gene.

Example 5 : The production of the first and second parent plants of this invention.

First parent plants and second parent plants (i.e. maintainer plants) according to the invention are produced along the lines set out in Figure 1.

The male-sterile plants of step 1 are those produced in Example 1. The corn plants transformed with the color-linked restorer gene of step 2 are those produced in Example 4.

A plant of Example 1 and a plant of Example 4 are crossed (Step 3) and the progeny plants with the genotype S/-, Rf/- are selected (Step 4), e.g. by demonstrating

the presence of both the S and Rf genes in the nuclear genome (e.g. by means of PCR).

The plants selected in Step 4 are then crossed with the male-sterile plants with genotype S/- (Step 5). The colored seeds (i.e. those containing the Rf gene) are selected, grown into plants, and examined for the presence of both the S and Rf genes (e.g. by PCR). The plants containing both the S and Rf genes are selfed and the seeds of each plant are examined on seed color (red or yellow). From the progeny of the selfings the non-colored seeds are grown into plants (step 6). The progeny of the selfings in which all noncolored seeds grow into male-sterile plants are retained (Step 6). These male-sterile plants are all homozygous for the S gene and are crossed with their fertile siblings (of genotype S/S,Rf/Rf or S/S,Rf/-) (Step 7). For some crossings the seeds harvested from the male-sterile plants are 50% colored and 50% non-colored (step 7). The colored seeds all grow into fertile corn plants of genotype S/S,Rf/- which are the maintainer plants, or the second parent plants, of the present invention. The noncolored seeds all grow into male-sterile plants of the genotype S/S,-/- which are the first parent plants of this invention (Step 7).

The first and second parent plants are crossed and the seeds harvested from the male-sterile plants are separated on the basis of color (Step 8). All colored

seeds grow again in second parent plants and all noncolored seeds grow in first parent plants, thereby establishing an easy maintenance of a pure male-sterile line of corn.

5

If the plant DNA that is flanking the S gene in the plants of Example 1 has been characterized, the progeny of the cross in Step 5 with genotypes S/S,-/- and S/S,R/- can be easily identified by means of PCR using probes corresponding to the flanking plant DNA. In this way Step 6 can be skipped because the plants of Step 5 which grow from colored seeds (genotype S/S,Rf/-) can be crossed directly to plants with genotype S/S,-/- (as in Step 7).

All publications cited in this application are hereby incorporated by reference.

Example 6 : Maintainer plants containing a color-linked restorer gene comprising the B-Peru coding region under control of the promoter of the Cl-S gene.

20

Using conventional techniques a chimeric gene is inserted between the EcoRI and HindIII sites of the polylinker of plasmid pUC19. The chimeric gene comprises the following elements in sequence:

25

i) the promoter region of the Cl-S gene, i.e. the DNA fragment with the sequence of SEQ ID No. 1 from nucleotide positions 447 up to 1076 but containing at

nucleotide positions 935-939 the sequence TTAGG instead of TGCAG.

ii) a single C nucleotide

5 iii) the coding region and 3'untranslated region of the B-peru gene, i.e. the DNA fragment with the sequence of SEQ ID No. 7 from nucleotide positions 576 up to 4137.

10 This plasmid (designated as pLH52), together with plasmid pCOL9S of Example 2 (comprising a C1-S gene) and pTS256 of SEQ ID No. 2 (comprising the following chimeric genes: P35S-bar-3'nos and PTA29-barstar-3'nos), is used to transform corn essentially as described in Example 4. The transformed plants are then used to obtain second parent plants as described in Example 5.

15

Example 7: Maintainer plants containing a color-linked restorer gene comprising the B-Peru coding region under control of the 35S promoter.

20 Using conventional techniques a chimeric gene is inserted between the EcoRI and HindIII sites of the polylinker of plasmid pUC19. the chimeric gene comprises the following elements in sequence:

25 i) The promoter region of the 35S promoter, i.e. the DNA fragment of pDE110 which essentially has the sequence as described in SEQ ID No. 4 of WO 92/09696 (which is incorporated herein by reference) from nucleotide positions 396 up to 1779

ii) the coding region and 3'untranslated region of the B-peru gene, i.e. the DNA fragment with the sequence of SEQ ID No. 7 from nucleotide positions 576 up to 4137.

5 This plasmid (designated as pP35S-Bp), together with plasmid pCOL9S of Example 2 (comprising a C1-S gene) and pTS256 of SEQ ID No. 2 (comprising the following chimeric genes: P35S-bar-3'nos and PTA-29-barstar-3'nos), is used to transform corn essentially as described in Example 4.
10 The transformed plants are then used to obtain second parent plants as described in Example 5.

Alternatively plasmid p35SBperu as described in Goff et al, 1990, EMBO 9:2517-2522 is used instead of pP35SBp.

15

Example 8: Maintainer plants containing a color-linked restorer gene comprising the maize P gene coding region under the control of the promoter of the C1-S gene.

Using conventional techniques a chimeric gene is inserted
20 in the EcoR1 site of the polylinker of plasmid pUC19. The chimeric gene comprises the following elements in sequence:

i) the promoter region of the C1-S gene, i.e. the DNA fragment with the sequence of SEQ ID No. 1 for nucleotide
25 positions 447 up to 1076 but containing at nucleotide positions 935-939 the sequence TTAGG instead of TGCAG ;

ii) a single C nucleotide;

iii) a DNA sequence comprising the coding region and 3'end untranslated region of the maize P gene as described by Grotewold et al in 1991, PNAS 88:4587-4591 (nucleotides 320-1517). The maize P gene is an anthocyanin regulatory gene which specifies red phlobaphene pigmentation, a flavonoid pigment involved in the biosynthetic pathway of anthocyanin. In fact, the protein encoded by the P gene activates, among others, the A1 gene required for both anthocyanin and phlobaphene pigmentation. Two cDNA clones have been isolated and sequenced by Grotewold et al and are described in the publication referred to above. It is the longer cDNA which is of particular interest for construction of this chimeric gene. However, alternatively, the coding region of the shorter transcript can also be used in this chimeric gene, as well as the P gene leader sequence instead of the CI-S gene leader sequence. The P gene does not require a functional R or B gene to produce pigmentation. The visible pigment that is produced in the seeds of the maintainer plants is phlobaphene, a flavonoid pigment (like anthocyanin) directly involved in anthocyanin biosynthesis.

iv) a DNA fragment containing the polyadenylation signal of the nopaline synthase gene of Agrobacterium tumefaciens, i.e. the DNA fragment with the sequence of

SEQ ID. No. 2 from nucleotide position 1600 up to nucleotide position 2909.

5 The resulting plasmid (designated as pPCS1-P), together with pTS256 of SEQ ID No. 2 is used to transform corn essentially as described in example 4. The transformed plants are then used to obtain second parent plants as described in example 5.

10 Example 9: Maintainer plants containing a color-linked restorer gene comprising the B-peru coding region under the control of the B-peru promoter.

Using conventional techniques a chimeric gene is inserted between the EcoR1 and the HindIII sites of the polylinker of plasmid pUC19. The chimeric gene comprises the
15 following elements in sequence:

- i) the promoter of the B-peru gene, i.e. a 1952 bp DNA sequence as disclosed in the EMBL databank under accession number X70791;
- 20 ii) the coding region and 3'untranslated region of the B-peru gene, i.e. the DNA fragment with the sequence of SEQ ID No. 7 from nucleotide position 576 up to 4137. This plasmid (designated aspCOL11), together with plasmid pCOL 9S of example 2 (comprising a C1-S gene) and pTS256
25 of SEQ ID No. 2 (comprising the following chimeric genes: P35S-bar-3'nos and PTA29-barstar-3'nos) is used to transform corn essentially as described in example 4. The

transformed plants are then used to obtain second parent plants as described in example 5.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: PLANT GENETIC SYSTEMS N.V.
- (B) STREET: Jozef Plateaustraat 22
- (C) CITY: Ghent
- (E) COUNTRY: Belgium
- (F) POSTAL CODE (ZIP): 9000
- (G) TELEPHONE: 32 9 235 84 11
- (H) TELEFAX: 32 9 224 06 94
- (I) TELEX: 11.361 Pgsgen

(ii) TITLE OF INVENTION: Use of anthocyanin genes to maintain male-sterile plants

(iii) NUMBER OF SEQUENCES: 7

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 08/254,776
- (B) FILING DATE: 06-JUN-1994

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4059 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: C1 gene of Zea mays

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 279..284
- (D) OTHER INFORMATION: /label= HpaI

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 447..452
- (D) OTHER INFORMATION: /label= EcoRI

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 1735..1740
- (D) OTHER INFORMATION: /label= AatII

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 1505..1510
- (D) OTHER INFORMATION: /label= EcoRI

- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2081..2086
 (D) OTHER INFORMATION:/label= XhoI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2418..2430
 (D) OTHER INFORMATION:/label= SfiI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2669..2674
 (D) OTHER INFORMATION:/label= SnaBI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2634..2639
 (D) OTHER INFORMATION:/label= SnaBI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:3008..3013
 (D) OTHER INFORMATION:/label= HpaI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1..1077
 (D) OTHER INFORMATION:/label= PC1
 /note= "region containing promoter of C1 gene"
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1078..2134
 (D) OTHER INFORMATION:/label= C1
 /note= "coding region of C1 gene"
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2135..2430
 (D) OTHER INFORMATION:/label= 3'C1
 /note= "region containing polyadenylation signal of C1
gene"
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1033..1038
 (D) OTHER INFORMATION:/label= TATA-Box
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1061..1062
 (D) OTHER INFORMATION:/label= transcript-init
 /note= "transcription initiation site"
- (ix) FEATURE:
 (A) NAME/KEY: intron
 (B) LOCATION:1211..1299
- (ix) FEATURE:
 (A) NAME/KEY: intron
 (B) LOCATION:1430..1575

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:935..939

(D) OTHER INFORMATION://label= C1-S

/note= "TGCAG sequence (in C1 gene) which in the C1-S sequence is changed to TTAGG"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TATCAACCTC CTGTGTTATT TTTAGTGACG GTTTCTTAAA AAACACCACT AGAAATCGTA	60
TTTTTATAGG TGGTTCCTTA AGAAACTGC ATGCAGAAAT CCATGACGGT TTTCTTAAGG	120
AACCGTATGT AGAAATACGA TTTCTAGTGA CGATCTTCTT AAGGAAACCA CCACTAAAAA	180
TTATTTTTAT CCTTAATTTT CGAGTTTTTC AAACGATCTC GTATGATGAA ACCATCAAAA	240
TAAAAGTTGT ACATCTCTAA AAGTTATGAA AATTTGTAGT TAACAACTTT TTTATTTGAA	300
CTCATTTTGG TTCTCAAAAA TTGCATCTAA ATTTGTCAAA TTTAAAATTC AAATTTTCCA	360
AACGACCTCG GATGAAAAAA GTGTCAAAAT GAAAGTTGTA GAACTTCAAA AGTTATTCAA	420
CTTTGTAGTC GACTATCTTT TTATTTGAAT TCGCTTACGG TCTCAAACAA GCAATTTACA	480
CTCAGTTGGT TGTAATATGT GGACAATAAA ACTACAACT AGACACAAAT CATACCATAG	540
ACGGAGTGGT AGCAGAGGGT ACGCGCGAGG GTGAGATAGA GGATTCTCCT AAAATAAATG	600
CACTTTAGAT GGGTAGGGTG GGGTGAGGCC TCTCCTAAAA TGAACTCGT TTAATGTTTC	660
TAAAAATAGT TTTCACCTGGT GATCCTTAGT TACTGGCATG TAAAAATGAT GATTTCTACT	720
GTCTCTCATA TGGACGGTTA TAAAAAATAC CATTATATTG AAAATAGGTC TCTGCTGCTA	780
CACTCGCCCT CATAGCAGAT CATGCATGCA CGCATCATTC GATCAGTTTT CGTTCTGATG	840
CAGTTTTCGA TAAATGCCAA TTTTTTAACT GCATACGTTG CCCTTGCTCA GCACCAGCAC	900
AGCAGTGTGCG TGTCGTCCAT GCATGCACTT TAGGTGCAGT GCAGGGCCTC AACTCGGCCA	960
CGTAGTTAGC GCCACTGCTA CAGATCGAGG CACCGGTCAG CCGGCCACGC ACGTCGACCG	1020
CGCGCGTGCA TTAAATACG CCGACGACGG AGCTTGATCG ACGAGAGAGC GAGCGCGATG	1080
GGGAGGAGGG CGTGTTCGCG GAAGGAAGGC GTTAAGAGAG GGGCGTGGAC GAGCAAGGAG	1140
GACGATGCCT TGGCCGCCTA CGTCAAGGCC CATGGCGAAG GCAAATGGAG GGAAGTGCCC	1200
CAGAAAGCCG GTAAACTAG CTAGTCTTTT TATTTCAATT TGGGATCATA TATATACCCC	1260
CGAGGCAAGA CCGGAGGACG ATCACGTGTG TGGGTGCAGG TTTGCGTCGG TGCGGCAAGA	1320
GCTGCCGGCT GCGGTGGCTG AACTACCTCC GGCCCAACAT CAGGCGCGGC AACATCTCCT	1380
ACGACGAGGA GGATCTCATC ATCCGCCTCC ACAGGCTCCT CGGCAACAGG TCTGTGCAGT	1440
GGCCAGTGGT GGGCTAGCTT ATTACACGAG CTGACGACGA GGCGATCGAT CGAGCGTCTG	1500
CTGCGAATTC ATCTGTTCCG GTGTCGGCCG TGTGAGAGTG AGCTCATTCA TATGTACATG	1560

CGTGTTGGCG CGCAGGTGGT CGCTGATTGC AGGCAGGCTG CCTGGCCGAA CAGACAATGA	1620
AATCAAGAAC TACTGGAACA GCACGCTGGG CCGGAGGGCA GGC GCCGCG CCGGCGCCGG	1680
CGGCAGCTGG GTCGTCGTCG CGCCGGACAC CGGCTCGCAC GCCACCCCGG CCGCGACGTC	1740
GGGCGCCTGC GAGACCGGCC AGAATAGCGC CGCTCATCGC GCGGACCCCG ACTCAGCCGG	1800
GACGACGACG ACCTCGGCGG CGGCGGTGTG GGC GCCAAG GCCGTGCGGT GCACGGGCGG	1860
ACTCTTCTTC TTCCACCGGG ACACGACGCC GGC GCACGCG GGC GAGACGG CGACGCCAAT	1920
GGCCGGTGGA GGTGGAGGAG GAGGAGGAGA AGCAGGGTCG TCGGACGACT GCAGCTCGGC	1980
GGCGTCGGTA TCGCTTCGCG TCGGAAGCCA CGACGAGCCG TGCTTCTCCG GCGACGGTGA	2040
CGGCGACTGG ATGGACGACG TGAGGGCCCT GGCCTCGTTT CTCGAGTCCG ACGAGGACTG	2100
GCTCCGCTGT CAGACGGCCG GGCAGCTTGC GTAGACAACA AGTACACGTA TAGATGTCCA	2160
ATAAGCACGA GGCCCGCGAG CCCGGCACGA AGCCCGCTTT TTGGGCCCGG TCCGAGCCCG	2220
GCACGGCCCG GTTATATGCA GACCCGGGCC GGC CGGCAC GAATAAGCGG GCCGGGCTCG	2280
GACAGGAAAT TAGGCACGGT GAGCTAGCCC GGCACGGCCC GTTTAGGTCT AAGCCCGTTA	2340
AGCCCGTTTT TTTACACTAA AACGTGCTTC TCGGCCGCA TAGCCCGCTT CTCGGCCCGC	2400
TTTTTTCGTG CTAAACGGGC CGGCCCGGCC CGGTTTAGGC CCGTTGCGGG CCGGGCTCGG	2460
ACAGGAAATT GAGCCCGCGT GCTTAGCCGG CCCGGCCCGG TTTTTTAATC GTGCCTGGCG	2520
GGCCAGGCCC AAAACGGGCC GGGCTTACC GGGCCCGGC CGGACCGGC CGGGCGGCCC	2580
GTTTGGACAT CTCTAAGTAC ACGTATGGAG GAGAATATAT ATATAGTCAT GCGTACGTAT	2640
AGATTTTTTC ATCCGATCCC AACAGAAATA CGTATGAAAA TGCTCTTCGT TCTTTTTTCAT	2700
TTATCATATC TATACTATAC TTAAAACACC AGTTTCAACG GTCGTCATGC GTCATTTTTT	2760
TACAAATAAC CCCTCACAGC TATTTCAAAT TAATCCGCTG CACGTCTATA GATGCCAAAC	2820
GACGCCCCA ACGGGCTAGA TGCACGCGG CCACAACAT GGCACAGGCA CGTCATGCCG	2880
GCCTGCTAAC TGTGTCGGGC TAGCCCGTTA GCCCGTCGAT CCATTTAATT AAATTAGCGT	2940
AACGACGCCC GACACGGGCT AGATGCACGT GGGCCACAAC TATGGCACAT GCACGTCATG	3000
CCGGCCTGTT AACTGTGTCG GGCCAGTCTG TTAGCCCAT TATCCATTTA ATTAAATCAG	3060
CGTAAAATGT TAAAAACGGT GCAGGAGGTG GGGTTCGAAC CCATACCCTG ATGGAAGAAG	3120
GGCGGGAGAC ACTGGGTGAA ACTGTCTAAC CAGTAGAATA TCTATCACGC TAAGATGTTT	3180
TTAATATTGA ATATAATTG TATATAAGCA TATAAGTTTT TTTGTAAAAT AAAAAATAAT	3240
CGTGTCGGGC CGGGCCATCA CTACTGGCCG AGGCTACAAC CCAAGCACGA CACGACGTTT	3300
TTGGCTCTTG CAAGCATTAG GTCGTTTCTG AGACCATATT GGC GCAATGG ACTACATGAT	3360
GTTTGGGGTT GCTGAATTGA ATGGAGCAGC AATAATTTGT CACACTAACA GCAAAATGAA	3420
AGGTTATTTG TTGGTTTTAA ACGTTAGTAA TTGCTACGAA GTAGCATAAT TTATATGGAG	3480

```

CGCATCCAGT TTTTATTGAT GCCTGACTTT AGCAATCACT CCATATTTTG ATCTATCTTT 3540
TTTATAAGTT TGACTTCATG GGACTTATTT TAGAACTTGA TCTCACAAAC TTTCTCTTAT 3600
TTTGTCTCTA TATGATGAAA TTGTGTCATT TTATAATCTT TGTTCAATCA GTCAATCGTT 3660
GTGAACTCTC TTCTAATCAC TCACTTCATT AGTTGTGTTG TACCAAGACA TATTTGCATA 3720
GAGTAAACAA TAACATCAGT TAGCCAAATC AAAAAATATA TTATACAGAG AGCGGAGACA 3780
ATCAAATAAA AAATCTTGAA ATTTTTTTTAA TGGATAGTTT ACGTGGGTAT TGTTGTAAGC 3840
CGTCGCAACG CACGGGCAAC CGACTAGTTT TAGTTTATAA ATTAATAAAC GTACGACAAA 3900
TATTAAGAAC GCCACCTTTC CATGCCTACG CGCGCGTGAG ACACGACCGG GGCACGTCAG 3960
ACGTGTGCCC CTGTTGTATA ATTTATTTAC TTTTAAATGA CTATGTGCTG TTGTTGCCG 4020
TTGGCTTCAT CGTGTTTCGTA GCCATGCATA AATCCAGCG 4059

```

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4896 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: circular
- (ii) MOLECULE TYPE: DNA (genomic)
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: plasmid pTS256, linearized at HindIII
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: complement (39..317)
 - (D) OTHER INFORMATION: /label= 3'nos
/note= "3' regulatory sequence containing the polyadenylation signal of the nopaline synthase gene of Agrobacterium T-DNA"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: complement (318..869)
 - (D) OTHER INFORMATION: /label= bar
/note= "coding region of bar gene of Streptomyces hygroscopicus"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: complement (870..1702)
 - (D) OTHER INFORMATION: /label= P35S
/note= "35S promoter of Cauliflower Mosaic Virus"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 1740..2284
 - (D) OTHER INFORMATION: /label= PTA29
/note= "promoter of TA29 gene of Nicotiana tabacum"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION:2285..2557
- (D) OTHER INFORMATION:/label= barstar
/note= "coding region of barstar gene of Bacillus amyloliquefaciens"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION:2558..2879
- (D) OTHER INFORMATION:/label= 3'nos
/note= "3' regulatory sequence containing the polyadenylation signal of the nopaline synthase gene of Agrobacterium T-DNA"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION:1..38
- (D) OTHER INFORMATION:/label= pUC19
/note= "pUC19 derived sequence"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION:2880..4896
- (D) OTHER INFORMATION:/label= pUC19
/note= "pUC19 derived sequence"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION:3004..3009
- (D) OTHER INFORMATION:/label= EcoRI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

AGCTTGCATG CCTGCAGGTC GACTCTAGAG GATCTTCCCG ATCTAGTAAC ATAGATGACA	60
CCGCGCGCGA TAATTTATCC TAGTTTGCGC GCTATATTTT GTTTTCTATC GCGTATTAAA	120
TGTATAATTG CGGGACTCTA ATCATAAAAA CCCATCTCAT AAATAACGTC ATGCATTACA	180
TGTTAATTAT TACATGCTTA ACGTAATTCA ACAGAAATTA TATGATAATC ATCGCAAGAC	240
CGGCAACAGG ATTCAATCTT AAGAACTTT ATTGCCAAAT GTTTGAACGA TCTGCTTCGG	300
ATCCTAGACG CGTGAGATCA GATCTCGGTG ACGGGCAGGA CCGGACGGGG CGGTACCGGC	360
AGGCTGAAGT CCAGCTGCCA GAAACCCAGC TCATGCCAGT TCCCGTGCTT GAAGCCGGCC	420
GCCCGCAGCA TGCCGCGGGG GGCATATCCG AGCGCCTCGT GCATGCGCAC GCTCGGGTCG	480
TTGGGCAGCC CGATGACAGC GACCACGCTC TTGAAGCCCT GTGCCTCCAG GGAATTCAGC	540
AGGTGGGTGT AGAGCGTGGA GCCCAGTCCC GTCCGCTGGT GGCGGGGGGA GACGTACACG	600
GTCGACTCGG CCGTCCAGTC GTAGGCGTTG CGTGCCTTCC AGGGGCCCCG GTAGGCGATG	660
CCGGCGACCT CGCCGTCCAC CTCGGCGACG AGCCAGGGAT AGCGCTCCCG CAGACGGACG	720
AGGTCGTCCG TCCACTCCTG CGGTTCTGCG GGCTCGGTAC GGAAGTTGAC CGTGCTTGTC	780
TCGATGTAGT GGTGACGAT GGTGCAGACC GCCGGCATGT CCGCCTCGGT GGCACGGCGG	840
ATGTCGGCCG GCGTCGTTT TGGGTCCATG GTTATAGAGA GAGAGATAGA TTTATAGAGA	900

GAGACTGGTG	ATTTACGCGT	GTCCTCTCCA	AATGAAATGA	ACTTCCTTAT	ATAGAGGAAG	960
GGTCTTGCGA	AGGATAGTGG	GATTGTGCGT	CATCCCTTAC	GTCAGTGGAG	ATGTCACATC	1020
AATCCACTTG	CTTTGAAGAC	GTGGTTGGAA	CGTCTTCTTT	TTCCACGATG	CTCCTCGTGG	1080
GTGGGGGTCC	ATCTTTGGGA	CCACTGTCGG	CAGAGGCATC	TTGAATGATA	GCCTTTCCTT	1140
TATCGCAATG	ATGGCATTTG	TAGGAGCCAC	CTTCCTTTTC	TACTGTCCTT	TCGATGAAGT	1200
GACAGATAGC	TGGGCAATGG	AATCCGAGGA	GGTTTCCCGA	AATTATCCTT	TGTTGAAAAG	1260
TCTCAATAGC	CCTTTGGTCT	TCTGAGACTG	TATCTTTGAC	ATTTTGGAG	TAGACCAGAG	1320
TGTCGTGCTC	CACCATGTTG	ACGAAGATTT	TCTTCTTGTC	ATTGAGTCGT	AAAAGACTCT	1380
GTATGAACTG	TTGCCAGTC	TTCACGGCGA	GTTCTGTTAG	ATCCTCGATT	TGAATCTTAG	1440
ACTCCATGCA	TGGCCTTAGA	TTCAGTAGGA	ACTACCTTTT	TAGAGACTCC	AATCTCTATT	1500
ACTTGCCTTG	GTTTATGAAG	CAAGCCTTGA	ATCGTCCATA	CTGGAATAGT	ACTTCTGATC	1560
TTGAGAAATA	TGTCTTTCTC	TGTGTTCTTG	ATGCAATTAG	TCCTGAATCT	TTTGACTGCA	1620
TCTTTAACCT	TCTTGGGAAG	GTATTTGATC	TCCTGGAGAT	TGTTACTCGG	GTAGATCGTC	1680
TTGATGAGAC	CTGCTGCGTA	GGAGCTTGCA	TGCCTGCAGG	TCGACTCTAG	AGGATCCCCA	1740
TCTAGCTAAG	TATAACTGGA	TAATTTGCAT	TAACAGATTG	AATATAGTGC	CAAACAAGAA	1800
GGGACAATTG	ACTTGTCACT	TTATGAAAGA	TGATTCAAAC	ATGATTTTTT	ATGTACTAAT	1860
ATATACATCC	TACTCGAATT	AAAGCGACAT	AGGCTCGAAG	TATGCACATT	TAGCAATGTA	1920
AATTAAATCA	GTTTTTGAAT	CAAGCTAAAA	GCAGACTTGC	ATAAGGTGGG	TGGCTGGACT	1980
AGAATAAACA	TCTTCTCTAG	CACAGCTTCA	TAATGTAATT	TCCATAACTG	AAATCAGGGT	2040
GAGACAAAAT	TTTGGTACTT	TTTCCTCACA	CTAAGTCCAT	GTTTGCAACA	AATTAATACA	2100
TGAAACCTTA	ATGTTACCCT	CAGATTAGCC	TGCTACTCCC	CATTTTCCTC	GAAATGCTCC	2160
AACAAAAGTT	AGTTTTGCAA	GTTGTTGTGT	ATGTCTTGTC	CTCTATATAT	GCCCTTGTGG	2220
TGCAAGTGTA	ACAGTACAAC	ATCATCACTC	AAATCAAAGT	TTTTACTTAA	AGAAATTAGC	2280
TACCATGAAA	AAAGCAGTCA	TTAACGGGGA	ACAAATCAGA	AGTATCAGCG	ACCTCCACCA	2340
GACATTGAAA	AAGGAGCTTG	CCCTTCCGGA	ATACTACGGT	GAAAACCTGG	ACGCTTTATG	2400
GGATTGTCTG	ACCGGATGGG	TGGAGTACCC	GCTCGTTTTG	GAATGGAGGC	AGTTTGAACA	2460
AAGCAAGCAG	CTGACTGAAA	ATGGCGCCGA	GAGTGTGCTT	CAGGTTTTC	GTGAAGCGAA	2520
AGCGGAAGGC	TGCGACATCA	CCATCATACT	TTCTTAATAC	GATCAATGGG	AGATGAACAA	2580
TATGGAAACA	CAAACCCGCA	AGCTTGGTCT	AGAGGATCCG	AAGCAGATCG	TTCAAACATT	2640
TGGCAATAAA	GTTTCTTAAG	ATTGAATCCT	GTTGCCGGTC	TTGCGATGAT	TATCATATAA	2700
TTTCTGTTGA	ATTACGTTAA	GCATGTAATA	ATTAACATGT	AATGCATGAC	GTTATTTATG	2760

AGATGGGTTT TTATGATTAG AGTCCCGCAA TTATACATTT AATACGCGAT AGAAAACAAA	2820
ATATAGCGCG CAAACTAGGA TAAATTATCG CGCGCGGTGT CATCTATGTT ACTAGATCGG	2880
GAAGATCCCC GGGTACCGAG CTCGAATTCT GATCAGGCCA ACGCGCGGGG AGAGGCGGTT	2940
TGCGTATTGG GCGCTCTTCC GCTTCCTCGC TCACTGACTC GCTGCGCTCG GTCGTTCCGC	3000
TGCGGCGAGC GGTATCAGCT CACTCAAAGG CGGTAATACG GTTATCCACA GAATCAGGGG	3060
ATAACGCAGG AAAGAACATG TGAGCAAAAG GCCAGCAAAA GGCCAGGAAC CGTAAAAAGG	3120
CCGCGTTGCT GCGCTTTTTT CATAGGCTCC GCCCCCTGA CGAGCATCAC AAAAATCGAC	3180
GCTCAAGTCA GAGGTGGCGA AACCCGACAG GACTATAAAG ATACCAGGCG TTTCCCCCTG	3240
GAAGCTCCCT CGTGCGCTCT CCTGTTCCGA CCCTGCCGCT TACCGGATAC CTGTCCGCCT	3300
TTCTCCCTTC GGGAAGCGTG GCGCTTTCTC AATGCTCAG CTGTAGGTAT CTCAGTTCGG	3360
TGTAGGTCGT TCGCTCCAAG CTGGGCTGTG TGCACGAACC CCCCCTCAG CCCGACCGCT	3420
GCGCCTTATC CGGTAACAT CGTCTTGAGT CCAACCCGGT AAGACACGAC TTATCGCCAC	3480
TGGCAGCAGC CACTGGTAAC AGGATTAGCA GAGCGAGGTA TGTAGGCGGT GCTACAGAGT	3540
TCTTGAAGTG GTGGCCTAAC TACGGCTACA CTAGAAGGAC AGTATTTGGT ATCTGCGCTC	3600
TGCTGAAGCC AGTTACCTTC GGAAAAAGAG TTGGTAGCTC TTGATCCGGC AAACAAACCA	3660
CCGCTGGTAG CGGTGGTTTT TTTGTTTGCA AGCAGCAGAT TACGCGCAGA AAAAAGGAT	3720
CTCAAGAAGA TCCTTTGATC TTTTCTACGG GGTCTGACGC TCAGTGAAC GAAAACTCAC	3780
GTTAAGGGAT TTTGGTCATG AGACTCGAGC CAAAAAGGAT CTTACCTAG ATCCTTTTAA	3840
ATTAATAATG AAGTTTTAAA TCAATCTAAA GTATATATGA GTAACTTGG TCTGACAGTT	3900
ACCAATGCTT AATCAGTGAG GCACCTATCT CAGCGATCTG TCTATTTCTG TCATCCATAG	3960
TTGCCTGACT CCCCCTCGTG TAGATAACTA CGATACGGGA GGGCTTACCA TCTGGCCCCA	4020
GTGCTGCAAT GATACCGCGA GACCCACGCT CACCGGCTCC AGATTTATCA GCAATAAACC	4080
AGCCAGCCGG AAGGGCCGAG CGCAGAAGTG GTCCTGCAAC TTTATCCGCC TCCATCCAGT	4140
CTATTAATTG TTGCCGGGAA GCTAGAGTAA GTAGTTCGCC AGTTAATAGT TTGCGCAACG	4200
TTGTTGCCAT TGCTACAGGC ATCGTGGTGT CACGCTCGTC GTTTGGTATG GCTTCATTCA	4260
GCTCCGGTTC CCAACGATCA AGGCGAGTTA CATGATCCCC CATGTTGTGC AAAAAAGCGG	4320
TTAGCTCCTT CGGTCCTCCG ATCGTTGTCA GAAGTAAGTT GGCCGAGTG TTATCACTCA	4380
TGGTTATGGC AGCACTGCAT AATTCTCTTA CTGTCATGCC ATCCGTAAGA TGCTTTTCTG	4440
TGACTGGTGA GTAACAACC AAGTCATTCT GAGAATAGTG TATGCGGCGA CCGAGTTGCT	4500
CTTGCCCGGC GTCAATACGG GATAATACCG CGCCACATAG CAGAACTTTA AAAGTGCTCA	4560
TCATTGGAAA ACGTTCTTCG GGGCGAAAAC TCTCAAGGAT CTTACCGCTG TTGAGATCCA	4620
GTTGATGTA ACCCACTCGT GCACCAACT GATCTTCAGC ATCTTTTACT TTCACCAGCG	4680

TTTCTGGGTG AGCAAAAACA GGAAGGCAAA ATGCCGCAAA AAAGGGAATA AGGGCGACAC 4740
GGAAATGTTG AATACTCATA CTCTTCCTTT TTCAATATTA TTGAAGCATT TATCAGGGTT 4800
ATTGTCTCAT GAGCGGATAC ATATTTGAAT GTATTTAGAA AAATAAACAA ATAGGGGTTT 4860
CGCGCACATT TCCCCGAAAA GTGCCACCTG ACGTCA 4896

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 3544 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: EcoRI-HindIII region of plasmid pTS200
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION:3227..3504
 - (D) OTHER INFORMATION:/label= 3'nos
/note= "3' regulatory sequence containing the polyadenylation signal of the nopaline synthase gene of Agrobacterium T-DNA"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION:2675..3226
 - (D) OTHER INFORMATION:/label= bar
/note= "coding region of bar gene of Streptomyces hygroscopicus"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION:1841..2674
 - (D) OTHER INFORMATION:/label= P35S
/note= "35S promoter of Cauliflower Mosaic Virus"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION:complement (626..1803)
 - (D) OTHER INFORMATION:/label= PCA55
/note= "promoter of CA55 gene of Zea mays"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION:complement (353..625)
 - (D) OTHER INFORMATION:/label= barstar
/note= "coding region of barstar gene of Bacillus amyloliquefaciens"
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION:complement (30..352)
 - (D) OTHER INFORMATION:/label= 3'nos
/note= "3' regulatory sequence containing the polyadenylation signal of the nopaline synthase gene of Agrobacterium T-DNA"

(ix) FEATURE:

- (A) NAME/KEY: -
 (B) LOCATION:1..6
 (D) OTHER INFORMATION:/label= EcoRI

(ix) FEATURE:

- (A) NAME/KEY: -
 (B) LOCATION:3539..3544
 (D) OTHER INFORMATION:/label= HindIII

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GAATTCGAGC TCGGTACCCG GGGATCTTCC CGATCTAGTA ACATAGATGA CACCGCGCGC	60
GATAATTTAT CCTAGTTTGC GCGCTATATT TTGTTTTCTA TCGCGTATTA AATGTATAAT	120
TGCGGGACTC TAATCATAAA AACCCATCTC ATAAATAACG TCATGCATTA CATGTTAATT	180
ATTACATGCT TAACGTAATT CAACAGAAAT TATATGATAA TCATCGCAAG ACCGGCAACA	240
GGATTCAATC TTAAGAACT TTATTGCCAA ATGTTTGAAC GATCTGCTTC GGATCCTCTA	300
GACCAAGCTT GCGGGTTTGT GTTCCATAT TGTTTATCTC CCATTGATCG TATTAAGAAA	360
GTATGATGGT GATGTCGCAG CTTTCCGCTT TCGCTTCACG GAAAACCTGA AGCACACTCT	420
CGGCGCCATT TTCAGTCAGC TGCTTGCTTT GTTCAAACCTG CCTCCATTCC AAAACGAGCG	480
GGTACTCCAC CCATCCGGTC AGAGAATCCC ATAAAGCGTC CAGGTTTTCA CCGTAGTATT	540
CCGGAAGGGC AAGCTCCTTT TTCAATGTCT GGTGGAGGTC GCTGATACTT CTGATTTGTT	600
CCCCGTTAAT GACTGCTTTT TTCATGGCTG CAGCTAGTTA GCTCGATGTA TCTTCTGTAT	660
ATGCAGTGCA GCTTCTGCGT TTTGGCTGCT TTGAGCTGTG AAATCTCGCT TTCCAGTCCC	720
TGCGTGTTTT ATAGTGCTGT ACGTTCGTGA TCGTGAGCAA ACAGGGCGTG CCTCAACTAC	780
TGTTTGGTT GGGTGACAGG CGCCAACTAC GTGCTCGTAA CCGATCGAGT GAGCGTAATG	840
CAACATTTTT TCTTCTTCTC TCGCATTGGT TTCATCCAGC CAGGAGACCC GAATCGAATT	900
GAAATCACAA ATCTGAGGTA CAGTATTTTT ACGTACCGT TCGTTCGAAG GTCTTCGACA	960
GGTCAAGGTA ACAAATCAG TTTTAAATTG TTGTTTCAGA TCAAAGAAAA TTGAGATGAT	1020
CTGAAGGACT TGGACCTTCG TCCAATGAAA CACTTGGACT AATTAGAGGT GAATTGAAAG	1080
CAAGCAGATG CAACCGAAGG TGGTGAAAGT GGAGTTTCAG CATTGACGAC GAAAACCTTC	1140
GAACGGTATA AAAAGAAGC CGCAATTAAA CGAAGATTG CCAAAAAGAT GCATCAACCA	1200
AGGGAAGACG TGCATACATG TTTGATGAAA ACTCGTAAAA ACTGAAGTAC GATTCCCCAT	1260
TCCCCTCCTT TTCTCGTTT TTTTAACTGA AGCAAAGAAT TTGTATGTAT TCCCTCCATT	1320
CCATATTCTA GGAGGTTTTG GCTTTTCATA CCCTCCTCCA TTTCAAATTA TTTGTCATAC	1380
ATTGAAGATA TACACCATT TAATTTATAC TAAATTACAG CTTTATAGATA CATATATTTT	1440
ATTATACACT TAGATACGTA TTATATAAAA CACCTAATTT AAAATAAAAA ATTATATAAA	1500
AAGTGTATCT AAAAAATCAA AATACGACAT AATTTGAAAC GGAGGGGTAC TACTTATGCA	1560

AACCAATCGT GGTAACCCTA AACCCCTATAT GAATGAGGCC ATGATTGTAA TGCACCGTCT	1620
GATTAACCAA GATATCAATG GTCAAAGATA TACATGATAC ATCCAAGTCA CAGCGAAGGC	1680
AAATGTGACA ACAGTTTTTT TTACCAGAGG GACAAGGGAG AATATCTATT CAGATGTCAA	1740
GTTCCCGTAT CACACTGCCA GGTCTTACT CCAGACCATC TTCCGGCTCT ATTGATGCAT	1800
ACCAGGAATT GATCTAGAGT CGACCTGCAG GCATGCAAGC TCCTACGCAG CAGGTCTCAT	1860
CAAGACGATC TACCCGAGTA ACAATCTCCA GGAGATCAAA TACCTTCCA AGAAGGTAA	1920
AGATGCAGTC AAAAGATTCA GGAATAATTG CATCAAGAAC ACAGAGAAAG ACATATTTCT	1980
CAAGATCAGA AGTACTATTC CAGTATGGAC GATTCAAGGC TTGCTTCATA AACCAAGGCA	2040
AGTAATAGAG ATTGGAGTCT CTAAAAAGGT AGTTCCTACT GAATCTAAGG CCATGCATGG	2100
AGTCTAAGAT TCAAATCGAG GATCTAACAG AACTCGCCGT GAAGACTGGC GAACAGTTCA	2160
TACAGAGTCT TTTACGACTC AATGACAAGA AGAAAATCTT CGTCAACATG GTGGAGCACG	2220
ACACTCTGGT CTACTCCAAA AATGTCAAAG ATACAGTCTC AGAAGACCAA AGGGCTATTG	2280
AGACTTTTCA ACAAAGGATA ATTTCTGGGAA ACCTCCTCGG ATTCCATTGC CCAGCTATCT	2340
GTCACTTCAT CGAAAGGACA GTAGAAAAGG AAGGTGGCTC CTACAAATGC CATCATTGCG	2400
ATAAAGGAAA GGCTATCATT CAAGATGCCT CTGCCGACAG TGGTCCCAA GATGGACCCC	2460
CACCCACGAG GAGCATCGTG GAAAAAGAAG ACGTTCCAAC CACGTCTTCA AAGCAAGTGG	2520
ATTGATGTGA CATCTCCACT GACGTAAGGG ATGACGCACA ATCCCACTAT CCTTCGCAAG	2580
ACCCTTCCTC TATATAAGGA AGTTCATTTT ATTTGGAGAG GACACGCTGA AATCACCAGT	2640
CTCTCTCTAT AAATCTATCT CTCTCTCTAT AACCATGGAC CCAGAACGAC GCCCGGCCGA	2700
CATCCGCCGT GCCACCGAGG CGGACATGCC GCGGTCTGC ACCATCGTCA ACCACTACAT	2760
CGAGACAAGC ACGGTCAACT TCCGTACCGA GCCGCAGGAA CCGCAGGAGT GGACGGACGA	2820
CCTCGTCCGT CTGCGGGAGC GCTATCCCTG GCTCGTCGCC GAGGTGGACG GCGAGGTCGC	2880
CGGCATCGCC TACGCGGGCC CCTGGAAGGC ACGCAACGCC TACGACTGGA CGGCCGAGTC	2940
GACCGTGATC GTCTCCCCC GCCACCAGCG GACGGGACTG GGCTCCACGC TCTACACCCA	3000
CCTGCTGAAG TCCCTGGAGG CACAGGGCTT CAAGAGCGTG GTCGCTGTCA TCGGGCTGCC	3060
CAACGACCCG AGCGTGCGCA TGCACGAGGC GCTCGGATAT GCGCCCGCG GCATGCTGCG	3120
GGCGGCCGGC TTCAAGCACG GGAAGTGGCA TGACGTGGGT TTCTGGCAGC TGGACTTCAG	3180
CCTGCCGGTA CCGCCCCGTC CGGTCCTGCC CGTCACCGAG ATCTGATCTC ACGCGTCTAG	3240
GATCCGAAGC AGATCGTTCA AACATTTGGC AATAAAGTTT CTTAAGATTG AATCCTGTTG	3300
CCGGTCTTGC GATGATTATC ATATAATTTC TGTTGAATTA CGTTAAGCAT GTAATAATTA	3360
ACATGTAATG CATGACGTTA TTTATGAGAT GGGTTTTTAT GATTAGAGTC CCGCAATTAT	3420

ACATTTAATA CGCGATAGAA AACAAAATAT AGCGCGCAAA CTAGGATAAA TTATCGCGCG 3480
 CGGTGTCATC TATGTTACTA GATCGGGAAG ATCCTCTAGA GTCGACCTGC AGGCATGCAA 3540
 GCTT 3544

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: oligonucleotide 1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CGTTTCTCGA ATCCGACGAG G 21

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4824 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: plasmid pCOL9

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 396..401
- (D) OTHER INFORMATION: /label= EcoRI

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 2367..2379
- (D) OTHER INFORMATION: /label= SfiI

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 884..888
- (D) OTHER INFORMATION: /label= C1-S
 /note= "TGCAG (in C1) which in C1-S allele is replaced with TTAGG"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

TCGCGCGTTT CGGTGATGAC GGTGAAAACC TCTGACACAT GCAGTCCCG GAGACGGTCA 60
 CAGCTTGCT GTAAGCGGAT GCCGGGAGCA GACAAGCCCG TCAGGGCGCG TCAGCGGGTG 120
 TTGGCGGGTG TCGGGGCTGG CTTAACTATG CGGCATCAGA GCAGATTGTA CTGAGAGTGC 180

ACCATATGCG GTGTGAAATA CCGCACAGAT GCGTAAGGAG AAAATACCGC ATCAGGCGCC	240
ATTCGCCATT CAGGCTGCGC AACTGTTGGG AAGGGCGATC GGTGCGGGCC TCTTCGCTAT	300
TACGCCAGCT GGCAGAAAGG GGATGTGCTG CAAGGCGATT AAGTTGGGTA ACGCCAGGGT	360
TTTCCCAGTC ACGACGTTGT AAAACGACGG CCAGTGAATT CGCTTACGGT CTCAAACAAG	420
CAATTTACAC TCAGTTGGTT GTAATATGTG GACAATAAAA CTACAACTA GACACAAATC	480
ATACCATAGA CGGAGTGGTA GCAGAGGGTA CGCGCGAGGG TGAGATAGAG GATTCTCCTA	540
AAATAAATGC ACTTTAGATG GGTAGGGTGG GGTGAGGCCT CTCCTAAAAT GAAACTCGTT	600
TAATGTTTCT AAAAATAGTT TTCACTGGTG ATCCTTAGTT ACTGGCATGT AAAATGATG	660
ATTTCTACTG TCTCTCATAT GGACGGTTAT AAAAAATACC ATTATATTGA AAATAGGTCT	720
CTGCTGCTAC ACTCGCCCTC ATAGCAGATC ATGCATGCAC GCATCATTCG ATCAGTTTTC	780
GTTCTGATGC AGTTTTCGAT AAATGCCAAT TTTTAACTG CATACGTTGC CCTTGCTCAG	840
CACCAGCACA GCAGTGTCTG GTCGTCCATG CATGCACTTT AGGTGCAGTG CAGGGCCTCA	900
ACTCGGCCAC GTAGTTAGCG CCACTGCTAC AGATCGAGGC ACCGGTCAGC CGGCCACGCA	960
CGTCGACCGC GCGCGTGCAT TTAAATACGC CGACGACGGA GCTTGATCGA CGAGAGAGCG	1020
AGCGCGATGG GGAGGAGGGC GTGTTGCGCG AAGGAAGGCG TTAAGAGAGG GCGTGGACG	1080
AGCAAGGAGG ACGATGCCTT GGCCGCCTAC GTCAAGGCCC ATGGCGAAGG CAAATGGAGG	1140
GAAGTGCCCC AGAAAGCCGG TAAACTAGC TAGTCTTTTT ATTTCATTTT GGGATCATAT	1200
ATATACCCCC GAGGCAAGAC CGGAGGACGA TCACGTGTGT GGGTGCAGGT TTGCGTCGGT	1260
GCGGCAAGAG CTGCCGGCTG CCGTGGCTGA ACTACCTCCG GCCCAACATC AGGCGCGGCA	1320
ACATCTCCTA CGACGAGGAG GATCTCATCA TCCGCCTCCA CAGGCTCCTC GGCAACAGGT	1380
CTGTGCAGTG GCCAGTGGTG GGCTAGCTTA TTACACGAGC TGACGACGAG GCGATCGATC	1440
GAGCGTCTGC TGCGAATTCA TCTGTTCCGG TGTCGGCCGT GTGAGAGTGA GCTCATTCAT	1500
ATGTACATGC GTGTTGGCGC GCAGGTGGTC GCTGATTGCA GGCAGGCTGC CTGGCCGAAC	1560
AGACAATGAA ATCAAGAACT ACTGGAACAG CACGCTGGGC CGGAGGGCAG GCGCCGGCGC	1620
CGGCGCCGGC GGCAGCTGGG TCGTCGTCGC GCCGGACACC GGCTCGCAGC CCACCCCGGC	1680
CGCGACGTCG GGCGCCTGCG AGACCGGCCA GAATAGCGCC GCTCATCGCG CGGACCCGA	1740
CTCAGCCGGG ACGACGACGA CCTCGGCGGC GCGGTGTGG GCGCCCAAGG CCGTGCGGTG	1800
CACGGGCGGA CTCTTCTTCT TCCACCGGGA CACGACGCCG GCGCACGCGG GCGAGACGGC	1860
GACGCCAATG GCCGGTGGAG GTGGAGGAGG AGGAGGAGAA GCAGGGTCGT CGGACGACTG	1920
CAGCTCGGCG GCGTCGGTAT CGCTTCGCGT CGGAAGCCAC GACGAGCCGT GCTTCTCCGG	1980
CGACGGTGAC GGCGACTGGA TGGACGACGT GAGGGCCCTG GCGTCGTTTC TCGAGTCCGA	2040
CGAGGACTGG CTCCGCTGTC AGACGGCCGG GCAGCTTGCG TAGACAACAA GTACACGTAT	2100

AGATGTCCAA TAAGCACGAG GCCC GCGAGC CCGGCACGAA GCCC GCTTTT TGGGCCCGGT	2160
CCGAGCCCGG CACGGCCCGG TTATATGCAG ACCCGGGCCG GCCC GGCACG AATAAGCGGG	2220
CCGGGCTCGG ACAGGAAATT AGGCACGGTG AGCTAGCCCG GCACGGCCCG TTTAGGTCTA	2280
AGCCCGTTAA GCCCGTTTTT TTACTATAA ACGTGCTTCT CGGCCCGCAT AGCCCGCTTC	2340
TCGGCCCGCT TTTTTCGTGC TAAACGGGCC GGGCCGGCCC GGTTTAGGCC CGTTGCGGGC	2400
CGGGCTCGGA CAGGAAATTG AGCCCGCGTG CTTAGCCGGC CCGGCCCGGT TTTTAAATCG	2460
TGCTTGGCGG GCCAGGCCCA AAACGGGCCG GGCTTCACCG GGGCCGGGCC GGACCGGGCC	2520
GGCGGGCCCG TTGGACATC TCTAAGTACA CGTATGGAGG AGAATATATA TATAGTCATG	2580
CGTACAGCTT GCGTAATCA TGGTCATAGC TGTTTCCTGT GTGAAATTGT TATCCGCTCA	2640
CAATTCCACA CAACATACGA GCCGGAAGCA TAAAGTGTA AGCCTGGGGT GCCTAATGAG	2700
TGAGCTAACT CACATTAATT GCGTTGCGCT CACTGCCCCG TTTCCAGTCG GGAAACCTGT	2760
CGTGCCAGCT GCATTAATGA ATCGGCCAAC GCGCGGGGAG AGGCGGTTTG CGTATTGGGC	2820
GCTCTTCCGC TTCCTCGCTC ACTGACTCGC TGCCTCGGT CGTTCGGCTG CGGCGAGCGG	2880
TATCAGCTCA CTCAAAGGCG GTAATACGGT TATCCACAGA ATCAGGGGAT AACGCAGGAA	2940
AGAACATGTG AGCAAAAGGC CAGCAAAAGG CCAGGAACCG TAAAAAGGCC GCGTTGCTGG	3000
CGTTTTTCCA TAGGCTCCGC CCCCTGACG AGCATCACAA AAATCGACGC TCAAGTCAGA	3060
GGTGGCGAAA CCCGACAGGA CTATAAAGAT ACCAGGCGTT TCCCCCTGGA AGCTCCCTCG	3120
TGCGCTCTCC TGTTCCGACC CTGCGCTTA CCGGATACCT GTCCGCCTTT CTCCCTTCGG	3180
GAAGCGTGGC GCTTCTCAA TGCTCACGCT GTAGGTATCT CAGTTCGGTG TAGGTCGTTT	3240
GCTCCAAGCT GGGCTGTGTG CACGAACCCC CCGTTCAGCC CGACCGCTGC GCCTTATCCG	3300
GTAACATATC TCTTGAGTCC AACC CGGTAA GACACGACTT ATCGCCACTG GCAGCAGCCA	3360
CTGGTAACAG GATTAGCAGA GCGAGGTATG TAGGCGGTGC TACAGAGTTC TTGAAGTGGT	3420
GGCCTAACTA CGGCTACACT AGAAGGACAG TATTGGTAT CTGCGCTCTG CTGAAGCCAG	3480
TTACCTTCGG AAAAGAGTT GGTAGCTCTT GATCCGGCAA ACAAAACCACC GCTGGTAGCG	3540
GTGGTTTTTT TGTTTGCAAG CAGCAGATTA CGCGCAGAAA AAAAGGATCT CAAGAAGATC	3600
CTTTGATCTT TTCTACGGGG TCTGACGCTC AGTGAACGA AAATCACGT TAAGGGATTT	3660
TGGTCATGAG ATTATCAAAA AGGATCTTCA CCTAGATCCT TTAAATTAA AAATGAAGTT	3720
TTAAATCAAT CTAAAGTATA TATGAGTAAA CTTGGTCTGA CAGTTACCAA TGCTTAATCA	3780
GTGAGGCACC TATCTCAGCG ATCTGTCTAT TTCGTTTCATC CATAGTTGCC TGAATCCCCG	3840
TCGTGTAGAT AACTACGATA CGGGAGGGCT TACCATCTGG CCCCAGTGCT GCAATGATAC	3900
CGCGAGACCC ACGCTCACCG GCTCCAGATT TATCAGCAAT AAACCAGCCA GCCGGAAGGG	3960

```

CCGAGCGCAG AAGTGGTCCT GCAACTTTAT CCGCCTCCAT CCAGTCTATT AATTGTTGCC      4020
GGGAAGCTAG AGTAAGTAGT TCGCCAGTTA ATAGTTTGCG CAACGTTGTT GCCATTGCTA      4080
CAGGCATCGT GGTGTCACGC TCGTCGTTTG GTATGGCTTC ATTCAGCTCC GGTTCCTAAC      4140
GATCAAGGCG AGTTACATGA TCCCCATGT TGTGCAAAAA AGCGGTTAGC TCCTTCGGTC      4200
CTCCGATCGT TGTCAGAAGT AAGTTGGCCG CAGTGTATC ACTCATGGTT ATGGCAGCAC      4260
TGCATAATTC TCTTACTGTC ATGCCATCCG TAAGATGCTT TTCTGTGACT GGTGAGTACT      4320
CAACCAAGTC ATTCTGAGAA TAGTGTATGC GGCGACCGAG TTGCTCTTGC CCGGCGTCAA      4380
TACGGGATAA TACCGCGCCA CATAGCAGAA CTTTAAAAGT GTCATCATT GGAAAACGTT      4440
CTTCGGGGCG AAAACTCTCA AGGATCTTAC CGCTGTTGAG ATCCAGTTCG ATGTAACCCA      4500
CTCGTGACC CAACTGATCT TCAGCATCTT TTACTTTTAC CAGCGTTTCT GGGTGAGCAA      4560
AAACAGGAAG GCAAAATGCC GCAAAAAAGG GAATAAGGCG GACACGGAAA TGTTGAATAC      4620
TCATACTCTT CCTTTTTCAA TATTATTGAA GCATTTATCA GGGTTATTGT CTCATGAGCG      4680
GATACATATT TGAATGTATT TAGAAAAATA AACAAATAGG GGTTCGCGC ACATTTCCCC      4740
GAAAAGTGCC ACCTGACGTC TAAGAAACCA TTATTATCAT GACATTAACC TATAAAAATA      4800
GGCGTATCAC GAGGCCCTTT CGTC                                             4824

```

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3915 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: EcoRI-HindIII region of plasmid pCOL13

(ix) FEATURE:

- (A) NAME/KEY: prim_transcript
- (B) LOCATION:188

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION:188..212

(ix) FEATURE:

- (A) NAME/KEY: intron
- (B) LOCATION:213..556

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION:557..718

(ix) FEATURE:

- (A) NAME/KEY: intron
- (B) LOCATION:719..1224

- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION:1225..2770
 (D) OTHER INFORMATION:/codon_start= 2
 /note= "exon containing 3' end coding region of B-peru' gene"
- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION:576..718
- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION:1225..2770
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1268..2770
 (D) OTHER INFORMATION:/note= "3' end of B-peru coding
 region which is derived from cDNA"
- (ix) FEATURE:
 (A) NAME/KEY: 3'UTR
 (B) LOCATION:2771..3272
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:3273..3891
 (D) OTHER INFORMATION:/label= 3'region
 /note= "further 3' flanking region of B-peru gene. This region
 is only of approximate length and the sequence needs to be
 confirmed."
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1..6
 (D) OTHER INFORMATION:/label= EcoRI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:11..16
 (D) OTHER INFORMATION:/label= XbaI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:45..50
 (D) OTHER INFORMATION:/label= KpnI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:265..270
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:329..334
 (D) OTHER INFORMATION:/label= XbaI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:835..840
 (D) OTHER INFORMATION:/label= BamHI

- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1268..1273
 (D) OTHER INFORMATION:/label= MluI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2787..2792
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2883..2888
 (D) OTHER INFORMATION:/label= MunI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2827..2832
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:3892..3897
 (D) OTHER INFORMATION:/label= SalI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:3910..3915
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:3892..3915
 (D) OTHER INFORMATION:/label= polylinker
 /note= "part of polylinker of pUC19"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GAATTCAGGT TCTAGACTAT TCTTGTGGCC TCGGGCGGAT GGCGGGTACC CATGTCTTCG	60
TTAGGCTTAT CTGACCGTGG AGATGAAATC TAACGGCTCA TAGAAATTAA ACTAACGTGG	120
ACACTCTGTC CTTGCTGTTT TGCTCCCTGC TCTTTATATA TAGAATGCCT GCTTGCAATTG	180
CACCCGTACG TACAGCGTAG CGCGGAGTGG AGGTGAGCTC CTCCTCCGAT TCTTGCCTAA	240
TCTTTGGTCT TTGCACACGT ACGAAAGCTT TTTGCATTGT TTCGTTGCTT CTGGATGATC	300
AGTACTCTTA GATATTAAGC GATACCGATC TAGAATCGAG TTGTTGTA CTCTCTGTCC	360
CTTTTGTGCA GCTATAACTA GCTAGGTTCC TTCGCATAGA GCCTCTCTAC AGAGTACAGA	420
CTAGCTAGCA GTGTCAGACA CGAAATGGAA ATGGTCACTT CCAAATTGCA CGAGCTGGAA	480
TTATATACTC TTCTGATCTT CTTACCCGTC TCTTTATAGC GTGATATGCG TTTCTGGCTT	540
CTTGCTTACG TGAAGGATTA TTAGTAAGGC GCGTGATGGC GCTCTCAGCT TCCCCGGCTC	600
AGGAAGAACT GCTGCAGCCT GCTGGGAGGC CGTTGAGGAA GCAGCTTGCT GCAGCCGCGA	660
GGAGCATCAA CTGGAGCTAT GCCCTCTTCT GGTCCATTTT AAGCACTCAA CGACCTCGGT	720
AAATGGAAGT CCTGATAATC TATAATTTGT CTGGCAGTTT TCTACAATC TGGTGAATGA	780

TCGTCACCTTC GTTTCCTGA TACATACATA CATACATATG AAATAAAGAA AGTCGGATCC	840
CGTGATGCGA TTGTAGTTAT CGCTTTTCCG CAAAATGGTT GCTTTTGTAA TCTGCATTCC	900
TTTTTTTCCC ACATCTTCTT CCTTCTCGCG AGTAACGACA ACGCCACCCG CGCCGCCTGC	960
CGCCCATCGC CCCGCCCTGG CCGGCGAGAG CCTCAGCCTA TTACACCAGC GGCGACCTCT	1020
TTTCCCCTTC CTCTACCCGC CCTCGTGGCC GTGCTCTCCC CCGCTCTAAC CTGGTCTGGC	1080
CGCCTCCGCT GCCACCTGCT CCGGCGGCCT CACCCGCGTC TTTCTCGTCC CTACCCTCTC	1140
TGCTCTGGG CGCATCATCA TCTGATATTC TGATGCAAAT AAAAAAGGTA TACCATATAA	1200
GGACAACAGA AAATATGGTT GCAGGGTGCT GACGTGGACG GACGGGTCTT ACAATGGCGA	1260
GGTGAAGACG CGTAAGATCT CCCACTCCGT GGAGCTGACA GCCGACCAGC TGCTCATGCA	1320
GAGGAGCGAG CAGCTCCGGG AGCTCTACGA GGCCCTCCGG TCCGGCGAGT GCGACCGCCG	1380
CGGCGCGCGG CCGGTGGGCT CGCTGTCGCC GGAGGACCTC GGGGACACCG AGTGGTACTA	1440
CGTGATCTGC ATGACCTACG CCTTCCTGCC GGGCCAAGGC TTGCCC GGCA GGAGTTCCGC	1500
GAGCAACGAG CATGTCTGGC TGTGCAACGC GCACCTCGCC GGCAGCAAGG ACTTCCCCCG	1560
CGCGCTCCTG GCCAAGAGCG CGTCCATTCA GACAATCGTC TGCATCCCGC TCATGGGTGG	1620
CGTGCTTGAG CTTGGTACTA CTGATAAGGT GCCGGAGGAC CCGGACTTGG TCAGCCGAGC	1680
AACCGTAGCA TTCTGGGAGC CGCAATGTCC GACATACTCG AAAGAGCCGA GCTCCAACCC	1740
GTCAGCATAC GAAACCGGGG AAGCCGCATA CATAGTCGTG TTGGAGGACC TCGATCACAA	1800
TGCCATGGAC ATGGAGACGG TGACTGCCGC CGCCGGGAGA CACGGAACCG GACAGGAGCT	1860
AGGAGAAGTC GAGAGCCCGT CAAATGCAAG CCTGGAGCAC ATCACCAAGG GGATCGACGA	1920
GTTCTACAGC CTCTGCGAGG AAATGGACGT GCAGCCGCTA GAGGATGCCT GGATAATGGA	1980
CGGGTCTAAT TTCGAAGTCC CGTCGTCAGC GCTCCCGGTG GATGGCTCAA GCGCACCCGC	2040
TGATGGTTCT CGCGCGACAA GTTTCGTGGT TTGGACGAGG TCATCGCACT CCTGCTCGGG	2100
TGAAGCGGCG GTGCCGGTCA TCGAAGAGCC GCAGAAATTG CTGAAGAAAG CGTTGGCCCG	2160
CGGCGGTGCT TGGGCGAACA CGAACTGCGG TGGCGGGGGC ACGACGGTAA CAGCCCAGGA	2220
AAACGGCGCC AAGAACCACG TCATGTCAGA GCGAAAGCGC CGGGAGAAGC TCAACGAGAT	2280
GTTCTCTGTT CTCAAGTCGT TGGTTCCTC CATTACAAG GTGGACAAAG CATCCATCCT	2340
CGCCGAAACG ATAGCCTATC TAAAGGAGCT TCAACGAAG GTACAAGAAC TGGAATCCAG	2400
GAGGCAAGGT GGCAGTGGGT GTGTCAGCAA GAAAGTCTGT GTGGGCTCCA ACTCCAAGAG	2460
GAAGAGCCCA GAGTTCGCCG GTGGCGCGAA GGAGCACCCC TGGGTCTCTC CCATGGACGG	2520
CACCAGCAAC GTCACCGTCA CCGTCTCGGA CACGAACGTG CTCCTGGAGG TGCAATGCCG	2580
GTGGGAGAAG CTCCTGATGA CACGGGTGTT CGACGCCATC AAGAGCCTCC ATTTGGACGC	2640

TCTCTCGGTT CAGGCTTCGG CACCAGATGG CTTCATGAGG CTCAAGATAG GAGCTCAGTT	2700
TGCAGGCTCC GGC GCCGTCG TGCCCGGAAT GATCAGCCAA TCTCTCGTA AAGCTATAGG	2760
GAAGCGATGA AAGGGCGCTA CATGTGAAGC TTAATTAATG GAAGCAAAC TGTATTTCTT	2820
GTGCAAAAGC TTACTATATA TTTCTGCAA ACCTGGTGTG CCTTGTTTG ATTTTCAGTC	2880
GCCAAATTGTG CCTTTGTTTT TATCAAGTGA TGATCTACAC ATATATATAG GAATATTGTA	2940
AAAGAGCGAT GTCATAGGGT TTTTTTATTA CAAGGAACAA GTCTTTCACG TGCTGGCCTC	3000
ACAAATCCTA AGAGAAAATC TGCTCATTTT GATTGCGTTC CGCAACAACT CTGTAATCCA	3060
TATCCTATGT ATCCGATCAA CTAGTCGATA GCCTCCGTCC GCCACATCAT CATATATCTA	3120
TCTATGTGTG TCATCTGACA CATACTCCTC GCGTACTGTG CTGACATATG ATACTGACAC	3180
AGCATATATG CATGCACATC GTCACACGAC ATATATCTCG CTACTACACA GATATTGGAT	3240
ACGATACTAT ATAGCATCAT GCGTGCTGCG ATNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3300
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3360
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3420
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3480
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3540
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3600
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3660
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3720
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3780
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN	3840
NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NGTCGACCTG	3900
CAGGCATGCA AGCTT	3915

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4137 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: EcoRI-HindIII region of plasmid pCOL13

(ix) FEATURE:

- (A) NAME/KEY: prim_transcript
- (B) LOCATION: 188

(ix) FEATURE:

- (A) NAME/KEY: exon

- (B) LOCATION:188..212
- (ix) FEATURE:
 (A) NAME/KEY: intron
 (B) LOCATION:213..556
- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION:557..718
- (ix) FEATURE:
 (A) NAME/KEY: intron
 (B) LOCATION:719..1224
- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION:1226..2771
 (D) OTHER INFORMATION:/codon_start= 2
 /note= "exon containing 3' end coding region of B-peru gene.
 this exon continues up to the polyadenylation site."
- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION:576..718
- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION:1226..2771
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1269..2771
 (D) OTHER INFORMATION:/note= "fragment of B-peru coding
 region which is derived from cDNA"
- (ix) FEATURE:
 (A) NAME/KEY: 3'UTR
 (B) LOCATION:2772..4137
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1..6
 (D) OTHER INFORMATION:/label= EcoRI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:11..16
 (D) OTHER INFORMATION:/label= XbaI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:45..50
 (D) OTHER INFORMATION:/label= KpnI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:265..270
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:329..334
 (D) OTHER INFORMATION:/label= XbaI

- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:835..840
 (D) OTHER INFORMATION:/label= BamHI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:1269..1274
 (D) OTHER INFORMATION:/label= MluI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2788..2793
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2884..2889
 (D) OTHER INFORMATION:/label= MunI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:2828..2833
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:4114..4119
 (D) OTHER INFORMATION:/label= SalI
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:4132..4137
 (D) OTHER INFORMATION:/label= HindIII
- (ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION:4114..4137
 (D) OTHER INFORMATION:/label= polylinker
 /note= "part of polylinker of pUC19"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GAATTCAGGT TCTAGACTAT TCTTGTGGCC TCGGGCGGAT GGCGGGTACC CATGTCTTCG	60
TTAGGCTTAT CTGACCGTGG AGATGAAATC TAACGGCTCA TAGAAATTAA ACTAACGTGG	120
ACACTCTGTC CTTGCTGTTT TGCTCCCTGC TCTTTATATA TAGAATGCCT GCTTGCATTG	180
CACCCGTACG TACAGCGTAG CGCGGAGTGG AGGTGAGCTC CTCCTCCGAT TCTTGCCTAA	240
TCTTTGGTCT TTGCACACGT ACGAAAGCTT TTTGCATTGT TTCGTTGCTT CTGGATGATC	300
AGTACTCTTA GATATTAAGC GATACCGATC TAGAATCGAG TTGTTGTACT CTCTCTGTCC	360
CTTTTGTGCA GCTATAACTA GCTAGGTTCC TTCGCATAGA GCCTCTCTAC AGAGTACAGA	420
CTAGCTAGCA GTGTCAGACA CGAAATGGAA ATGGTCACTT CCAAATTGCA CGAGCTGGAA	480
TTATATACTC TTCTGATCTT CTTACCGTC TCTTTATAGC GTGATATGCG TTTCTGGCTT	540
CTTGCTTACG TGAAGGATTA TTAGTAAGGC GCGTGATGGC GCTCTCAGCT TCCCCGGCTC	600

AGGAAGAACT GCTGCAGCCT GCTGGGAGGC CGTTGAGGAA GCAGCTTGCT GCAGCCGCGA	660
GGAGCATCAA CTGGAGCTAT GCCCTCTTCT GGTCCATTTC AAGCACTCAA CGACCTCGGT	720
AAATGGAAGT CCTGATAATC TATAATTTGT CTGGCAGTTT TCTACAACTC TGGTGAATGA	780
TCGTCACTTC GTTTGCCTGA TACATACATA CATACATATG AAATAAAGAA AGTCGGATCC	840
CGTGATGCGA TTGTAGTTAT CGCTTTTCCG CAAAATGGTT GCTTTTGTAA TCTGCATTCTG	900
TTTTTTTCCC ACATCTTCTT CCTTCTCGCG AGTAACGACA ACGCCACC6C GCGCCGCGCTG	960
CCGCCCATCG CCCC6CCTTG GCCGGCGAGA GCCTCAGCCT ATTACACCAG CGGCGACCTC	1020
TTTTCCCCTT CCTCTCACCG CCCTCGTGGC CGTGCTCACC CCCGCTCTAA CCTGGTCTGG	1080
CCGCTCCGC TGCCACCTGC TCCGGCGGCC TCACCCGCGT CTTTCTCGTC CCTACCCTCT	1140
CTGCCTCTGG GCGCATCATC ATCTGATATT CTGATGCAAA GAAAAAGGT ATACCATATA	1200
AGGACAACAG AAAATATGGT TGCAGGGTGC TGACGTGGAC GGACGGGTTC TACAATGGCG	1260
AGGTGAAGAC GCGTAAGATC TCCCCTCCG TGGAGCTGAC AGCCGACCAG CTGCTCATGC	1320
AGAGGAGCGA GCAGCTCCGG GAGCTCTACG AGGCCCTCCG GTCCGGCGAG TCGGACCGCC	1380
GCGGCGCGCG GCCGGTGGGC TCGCTGTCGC CGGAGGACCT CGGGGACACC GAGTGGTACT	1440
ACGTGATCTG CATGACCTAC GCCTTCCTGC CGGGCCAAGG CTTGCCCCGC AGGAGTCCG	1500
CGAGCAACGA GCATGTCTGG CTGTGCAACG CGCACCTCGC CGGCAGCAAG GACTTCCCCC	1560
GCGCGCTCCT GGCCAAGAGC GCGTCCATTC AGACAATCGT CTGCATCCCG CTCATGGGTG	1620
GCGTGCTTGA GCTTGGTACT ACTGATAAGG TGCCGGAGGA CCCGGACTTG GTCAGCCGAG	1680
CAACCGTAGC ATTCTGGGAG CCGCAATGTC CGACATACTC GAAAGAGCCG AGCTCCAACC	1740
CGTCAGCATA CGAAACCGGG GAAGCCGCAT ACATAGTCGT GTTGGAGGAC CTCGATCACA	1800
ATGCCATGGA CATGGAGACG GTGACTGCCG CCGCCGGGAG ACACGGAACC GGACAGGAGC	1860
TAGGAGAAGT CGAGAGCCCG TCAAATGCAA GCCTGGAGCA CATCACCAG GGGATCGACG	1920
AGTTCTACAG CCTCTGCGAG GAAATGGACG TGCAGCCGCT AGAGGATGCC TGGATAATGG	1980
ACGGGTCTAA TTTCGAAGTC CCGTCGTCAG CGCTCCCGGT GGATGGCTCA AGCGCACCCG	2040
CTGATGGTTC TCGCGCGACA AGTTTCGTGG TTTGGACGAG GTCATCGCAC TCCTGCTCGG	2100
GTGAAGCGGC GGTGCCGGTC ATCGAAGAGC CGCAGAAATT GCTGAAGAAA GCGTTGGCCG	2160
GCGGCGGTGC TTGGGCGAAC ACGAACTGCG GTGGCGGGGG CACGACGGTA ACAGCCCAGG	2220
AAAACGGCGC CAAGAACCAC GTCATGTCAG AGCGAAAGCG CCGGGAGAAG CTCAACGAGA	2280
TGTTCTCGT TCTCAAGTCG TTGGTTCCCT CCATTCACAA GGTGGACAAA GCATCCATCC	2340
TCGCCGAAAC GATAGCCTAT CTAAAGGAGC TTCAACGAAG GGTACAAGAA CTGGAATCCA	2400
GGAGGCAAGG TGGCAGTGGG TGTGTCAGCA AGAAAGTCTG TGTGGGCTCC AACTCCAAGA	2460

GGAAGAGCCC AGAGTTCGCC GGTGGCGCGA AGGAGCACCC CTGGGTCCTC CCCATGGACG 2520
GCACCAGCAA CGTCACCGTC ACCGTCTCGG ACACGAACGT GCTCCTGGAG GTGCAATGCC 2580
GGTGGGAGAA GCTCCTGATG ACACGGGTGT TCGACGCCAT CAAGAGCCTC CATTGAGC 2640
CTCTCTCGGT TCAGGCTTCG GCACCAGATG GCTTCATGAG GCTCAAGATA GGAGCTCAGT 2700
TTGCAGGCTC CGGCGCCGTC GTGCCCAGAA TGATCAGCCA ATCTCTTCGT AAAGCTATAG 2760
GGAAGCGATG AAAGGGCGCT ACATGTGAAG CTTAATTAAT GGAAGCAAAC TTGTATTTCT 2820
TGTGCAAAAG CTTACTATAT ATTTCTGCAA AACCTGGTGT GCCTTGTTTT GATTTTCAGT 2880
CGCCAATTGT GCCTTTGTTT TTATCAAGTG ATGATCTACA CTATATATAT GGAATATTTG 2940
AAAAGAGCGA TGTCATAGGG TTTTTTTATT ACAAGGAACA AGTCTTTCAC GTGCTGGCCT 3000
CACAAATCCA AGAGAAAATC TGCTCATTTT GATTGGCTTC CGCAACAACCT CTGTAATCCA 3060
TATCCTTTGT ATCCGATCAA CTATGATACC TCCTCCCCCA TCTCTTTTTT TTTTATCTGC 3120
ACAATCTTCT ATTCTACTAT AATGAAACAA TAGAGCCACT ACCGAATATT TCCTCAAAAA 3180
TGTACAACAA ACTAGGGTGG TCCAAACAAA TGCCTAGAGG AGCTAGATTC TCTTAAATTA 3240
GACATCGGTT TCTTTTATCT CTTCCAGAAG GGATAAAAGT ATGTGTTTAT GGTCTTCAGT 3300
AATACATTGT TCGTTTCTCA TAGTCAATTT AGAGGTGTTT AAATGTACTT GAACTAATAG 3360
TTAGTTGGTT TAAAAATTAC TATTAAAATT AGTTAGTTAA TAAATAGCTA GCTAAATATT 3420
AGCTAATTTG TCAAAAGTAG CTAATAGCTG AATTATTAGC TATATTGTTT TGATGTCTTC 3480
AGCTAATTTT AGCAGATCAT TATTAGTTCT AGTGTATCTA AACACACCCT TAGTCAAAACA 3540
TGGTAAAAAA AAAGTTGATT CACTCATTGC TCATCGAAGA CGCAGATCAT GGCATCCCTC 3600
ACACGTTCTT CAGCCTACAC GGCACCTGCA TTGTAATTGC ATCTCATCTC ATCAACCCTT 3660
GTTGTGCATT ACTTGCCACA TCGGCCATCA ATTAACATTT TTTTGTCTCG TTCCTGAATT 3720
TCCTAACAAA TTTTCATCAA TGTACGCAGA GCTAAAGCTA GCTGTCGATG TCAGTTGACA 3780
GTTGACACCG ATGAATTTTA GAAAATTTAG TGTAAGTAC TATTTATAAT GTTCATGACA 3840
CCCATATAAA ATATGTTGAC ACCGGCAAAC CTTAAGGCTA GCTTCGCCCC TGCCATCAAC 3900
CTTACATCTA CATTCAACCAC GAGGTGTGCA CGGCCTAGGT TCGACTCCTA TGTATGCCT 3960
TGCTATCTAC AGATTCAGCA AGTGTGTGT TCCTTGTGT CACAATCTAC CTTTATTATA 4020
AAATTGATGT CATATCATGC CAAACAACAA ATAATTAATA TCGTGTGAAA TTTGAATTC 4080
TCTAACATGC TCAACCAACC TTACCCCTTC ACGGTCGACC TGCAGGCATG CAAGCTT 4137

SEQ. ID No. 7

actagtagcctgtcgcgcgcccattgcccgcgtggcggtgcttctcgccctggtaactgttctcgggcaaatga
ctattttccaagtaaacatattcaatgatttttgctattcttagcaaagtaatttcacttggacttttgtgc
caaaaacgcattggaaaaaatctccttggactccagcctaagggtgaaagtgtaaaaactgggaaaaatt
attgatgtttcgggcagttacttggctatgtaaattccataaccttttcaaaatatcctaaacattctttt
ctgtttctgcaacatacatgtttatcagttcttgaccttttgacgctacgaaagtccagttagtattcagg
ctttcgcaagtaaacctagaagtcacacggacattcatttttagcgtaccatgtcttttaggatgcactt
gttatcggtatgtctcctatgagacagaatgcacttgttatggtaactaaacaaaaaaataatttaatt
cgtgtgaaactttttcaaacctaccttccctgttcccgagggtccataataccagacacctaactcgcttg
cgcaatttagaagaaatcatgcatatacgtcaaggagggtgaaatatcaagcaaaaagaaagggtcat
cccacaaaagcccaaaactattgtagggaaaacacttgttttacctataattgagcgtcgatttgggtgtt
gctgatatttactgtcaaaccaagtcacatttaccagaatagtatctagaagaatccttttcacatcctc
tagcccgccaacatcctaccatttgacattgagaactaaaaaacaattgttccagacgaaagctaaag
tcgctttatacgattagctgcagtaggtgagcagcatctccgaacgctgggcatgacacgaccatgatag
acgacatggacattttgtcaaacacctgcatggcgctcaccagggaaaacaatccagcaggagaggtggga
gagagatggaaacaatttaattatgcaaacacggaggagacacaatttgaagagtgttcgtacacctacgg
caatcagcgaaacgatgagagagcataaccaagctcgggtcgtcagacacgaggagcggagcgtggcac
cgatggagatggagacagttgcgtgcgctttttgtggagggttctgttgggtgtcgggctcggcgaggc
ctgaacgcggtgggaagaagagcggcggtgggtgggaagaagagcagcgtcaggttcttagactattcttgtg
gcctcgggcggtatggcggttaccatgtcttctgttaggcttatctgacgctggagatgaaatctaacggc
tcatagaaattaaactaacgtggactcccagacgaaagctaaagtcgctttatacgattagctgcagtag
gtgagcacgatctccgaacgctgggcatgacacgaccatgatagacgacatggacattttgtcaaacacc
tgcattggcgctcaccagggaaaacaatccagcaggagaggtgggagagagatggaaacaatttaattatgca
aacacggaggagacacaatttgaagagtgttcgtacacctacggcaatcagcgaaacgatgagagagcat
accaagctcgggtcgtcagcaacgcggaggacggacgggtggcaccgatggagatggagacagttgcgtgc
cgttttttgtggagggttctgttgggtgtcgggcgtcggcgagcctgaacgcgggtgggaagaagagcttc
gtgttgggaagaagagcagacaggttcttagactattcttgtggcctcgggcggtatggcggttaccat
gtcttctgttaggcttatctgacgctggagatgaaatctaacggctcatagaaattaaactaacgtggaca
ctctgtccttgcgtgttttgcctcctgtctttatatatagaatgcctgottgcattgcacccgtacgtac
agcgtagcggcggtgaggtgagctcctcctccgatcttgcctaatotttgggtctttgcacacgtacg
aaagctttttgcattgtttcgttgccttctggatgatcagtagctttagatatttagcgataccgatctag
aatcgagtgtttgtactctctctgtcccttttgtgcagctataactagctaggttctcttgcataagagcc
tctctacagagtacagactagctagcagtgctcagacacgaaatggaaatgggtcacttccaaattgcacga
gctgggaattatatactcttctgatcttcttaccgctctctttatagcgtgatatgcgtttctggcttctt
gcttacgtgaaggattatttagtaaggcgcggtgatggcgctctcagcttccccggctcaggaagaactgct
gcagcctgctgggaggccgttgaggaagcagcttgctgcagccgcgaggagcatcaactggagctatgcc
ctcttctgggtccatttcaagcactcaacgacctcggtaaatggaaagtcctgataatctataatttgtctg
gcagttttctacaactctggtgaatgatcgtcacttgcgtttgcctgatacatatatacatatgaaa
taaagaaagtcggatcccgtagtgcatgttagttatcgcttttccgcaaaatgggttgccttttgaatct
gc1

CLAIMS

1. A plant consisting essentially of cells which comprise in their genome:

5

- a homozygous male-sterility genotype at a first genetic locus; and

- a color-linked restorer genotype at a second genetic locus, which is heterozygous (Rf/-) for a foreign DNA Rf

10

comprising:

a) a fertility-restorer gene capable of preventing the phenotypic expression of said male-sterility genotype, and

15

b) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of said plant which is capable of producing anthocyanin at least in the seeds of said plant, so that anthocyanin production in the seeds is visible externally.

20

2. The plant of claim 1 in which said color gene is capable of producing anthocyanin at least in the aleurone of the seeds of said plant.

25

3. The plant of claim 1, in which said first genetic locus is homozygous for a foreign RNA S (S/S) which comprises a male-sterility gene which when generated in cells of the plant renders the plant male-sterile without otherwise substantially affecting the growth and development of the plant.

4. The plant of claim 1, in which said first genetic locus is homozygous for a foreign DNA S (S/S) which comprises a male-sterility gene which comprises:

s1) a male-sterility DNA encoding a RNA, protein or polypeptide which, when produced or overproduced in a stamen cell of the plant, significantly disturbs the metabolism, functioning and/or development of said cell, and,

s2) a sterility promoter capable of directing expression of the male-sterility DNA selectively in the stamen cells, preferably the tapetum cells, of the plant; the male-sterility DNA being in the same transcriptional unit as, and under the control of, the sterility promoter,

and in which said fertility restorer gene in said second genetic locus comprises at least:

a1) a fertility-restorer DNA encoding a restorer RNA, protein or polypeptide which, when produced or overproduced in the same cell as said male-sterility

gene S, prevents the phenotypic expression of S,
and,

- 5 a2) a restorer promoter capable of directing expression
of the fertility-restorer DNA at least in the same
cells in which said male-sterility gene is
expressed, so that the phenotypic expression of said
male-sterility gene is prevented; the fertility-
restorer DNA being in the same transcriptional unit
as, and under the control of, the restorer promoter.

10

5. The plant of claim 1 in which said male-sterility DNA
encodes barnase and in which said fertility restorer DNA
encodes barstar.

15

6. The plant of claim 1 in which the sterility promoter
and/or the restorer promoter is selected from the group
consisting of PTA29, PCA55, PT72, PT42, and PE1.

20

7. The plant of claim 1 in which the homozygous male-
sterility genotype is endogenous and is homozygous for a
recessive allele m (m/m) and in which the fertility
restorer gene is the dominant allele M of said endogenous
male-sterility genotype.

25

8. The plant of claim 1, which is a cereal plant which is
selected from the group consisting of corn, wheat, and
rice.

9. The plant according to claim 1 wherein said anthocyanin regulatory gene is a shortened R, B or C1 gene or a combination of shortened R, B or C1 genes which
5 is functional for conditioning and regulating anthocyanin production in the aleurone.

10. The plant according to claim 9 wherein said anthocyanin regulatory gene is selected from the group
10 consisting of a shortened C1 or C1-S gene having a nucleotide sequence corresponding to the sequence between positions 447 and 2418 of SEQ ID No. 1, a shortened B-peru gene having a nucleotide sequence corresponding to the sequence between positions 1 and 3272 of SEQ of ID
15 NO. 6; and the Eco-SalI fragment having a length of about 4 ààà bp of pCOL13.

11. The plant according to claim 10 wherein said anthocyanin regularatory gene does not contain any
20 introns.

12. The plant according to claim 9 wherein said anthocyanin regulatory gene comprises a shortened C1 or C1-S gene and a shortened B-peru gene.
25

13. The plant according to claim 9 wherein said anthocyanin regulatory gene is a chimaric DNA comprising

a coding region of an R or B gene and/or C1 gene operably linked to an aleurone-specific promoter.

5 14. The plant according to claim 13 wherein said aleurone-specific promoter is selected from the group consisting of: the sequence between positions 1 to 1077 or 447 to 1077 of SEQ ID No. 1, and the sequence between positions 1-575 of sequence ID No. 6.

10 15. The plant according to claim 14, wherein said aleurone-specific promoter is selected from the group consisting of: the sequence between positions 1 to 1061 or 447 to 1061 of SEQ ID No. 1, and the sequence between positions 1 to 188 of SEQ ID No. 6.

15 16. A DNA comprising an anthocyanin regulatory gene which is a shortened R, B or C1 gene or a combination of shortened R, B or C1 genes which is functional for conditioning and regulating anthocyanin production in the
20 aleurone.

17. A DNA according to claim 16, which comprises a shortened C1 or C1-S gene and a shortened B-peru gene.

25 18. A DNA according to claim 16, which comprises at least one gene selected from the group consisting of a shortened B-peru gene having a nucleotide sequence

corresponding to the sequence between positions 1 and 3272 of SEQ ID No. 6, a shortened B-peru gene which is the EcoRI-SaII fragment with a length of about 4 000 bp of pCOL13 and the shortened C1 or C1-S gene having a
5 nucleotide sequence corresponding to the sequence between positions 447 and 2418 SEQ ID No. 1.

19. The DNA of claim 18 in which said shortened B-peru, C1 or C1-S gene is further characterized by not
10 containing any intron.

20. A DNA according to claim 16, wherein said shortened C1, C1-S or B-peru genes are operably linked to an aleurone-specific promoter selected from the group
15 consisting of: the sequence between positions 1 to 1077 or 447 to 1077 of SEQ ID No. 1 and the sequence between positions 1-575 of ID No. 6.

21. A DNA according to claim 19, wherein said aleurone-specific promoter is selected from the group consisting
20 of: the sequence between positions 1 to 1061 or 447 and 1061 of SEQ ID No. 1 and the sequence between positions 1 to 188 of SEQ. ID No. 6.

22. A DNA according to claim 16 which further comprises a
25 fertility-restorer gene capable of preventing the

phenotypic expression of a male-sterility genotype in a plant.

23. A DNA according to claim 22 wherein said fertility-restorer gene encodes barstar.

24. A DNA according to claim 23, wherein barstar is under the control of a promoter selected from the group consisting of PTA29, PCA55, PT72, PT42 and PE1.

25. An aleurone-specific promoter selected from the group consisting of: the sequence between positions 1 to 1077 or 447 to 1077 of SEQ ID No. 1 and the sequence between positions 1-575 of SEQ ID No. 6.

26. An aleurone-specific promoter selected from the group consisting of: the sequence between positions 1 to 1061 or 447 and 1061 of SEQ ID No. 1 and the sequence between positions 1 to 188 of SEQ ID No. 6.

27. A process to maintain a line of male-sterile plants, which comprises the following steps:

i) crossing

- a) a male-sterile parent plant of said line having in a first genetic locus, a homozygous male-sterility genotype and
- b) a maintainer parent plant of said line consisting essentially of cells which comprise, stably integrated in their nuclear genome:
- a homozygous male-sterility genotype at a first genetic locus; and
 - a colored-linked restorer genotype at a second genetic locus, which is heterozygous for a foreign DNA comprising:
 - i) a fertility-restorer gene capable of preventing the phenotypic expression of said male-sterility genotype, and
 - ii) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of said plant which is capable of producing anthocyanin at least in the seeds of said plant, so that anthocyanin production in the seeds is visible externally,
- ii) obtaining the seeds from said parent plants, and

iii) separating on the basis of color, the seeds in which no anthocyanin is produced and which grow into male-sterile parent plants.

5 28. A process according to claim 27, wherein the genome of said male-sterile parent plant does not contain at least one anthocyanin regulatory gene necessary for the regulation of anthocyanin biosynthesis in the seeds of said plant to produce externally visible anthocyanin in
10 said seeds.

29. The process of claim 28, wherein the genome of said male-sterile parent plant contains a first anthocyanin regulatory gene and the genome of said maintainer parent
15 plant contains a second anthocyanin regulatory gene which, when present with said first anthocyanin regulatory gene in the genome of a plant is capable of conditioning the production of externally visible anthocyanin in seeds.

20

30. A process to maintain a line of maintainer plants, which comprises the following steps:

i) crossing:

25

- a) a male-sterile parent plant of said line having, in a first genetic locus, a homozygous male-sterility genotype, and
- 5 b) a maintainer parent plant of said line consisting essentially of cells which comprise, stably integrated in their nuclear genome:
- a homozygous male-sterility genotype at a first genetic locus; and
 - 10 - a colored-linked restorer genotype at a second genetic locus, which is heterozygous for a foreign DNA comprising:
- 15 i) a fertility-restorer gene capable of preventing the phenotypic expression of said male-sterility genotype, and
 - 20 ii) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of said plant which is capable of producing anthocyanin at least in the seeds of said plant, so that anthocyanin production in the seeds is visible
 - 25 externally.

ii) obtaining the seeds from said male-sterile parent plant, and

iii) separating on the basis of color, the seeds in which anthocyanin is produced and which grow into maintainer parent plants.

31. A process according to claim 30, wherein the genome of said male-sterile parent plant does not contain at least one anthocyanin regulatory gene necessary for the regulation of anthocyanin biosynthesis in the seeds of said plant to produce externally visible anthocyanin in said seeds.

32. The process of claim 31, wherein the genome of said male-sterile parent plant contains a first anthocyanin regulatory gene and the genome of said maintainer parent plant contains a second anthocyanin regulatory gene which, when present with said first anthocyanin regulatory gene in the genome of a plant is capable of conditioning the production of externally visible anthocyanin in seeds.

33. A kit for maintaining a line of male-sterile or maintainer plants, said kit comprising:

a) a male-sterile parent plant of said line having, in a first genetic locus, a homozygous male-sterility genotype, and

5 b) a maintainer parent plant of said line consisting essentially of cells which comprise, integrated in their nuclear genome :

10 - a homozygous male-sterility genotype at a first genetic locus; and

 - a colored-linked restorer genotype at a second genetic locus, which is heterozygous for a foreign DNA comprising:

15

i) a fertility-restorer gene capable of preventing the phenotypic expression of said male-sterility genotype, and

20

ii) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of said plant which is capable of producing anthocyanin at least in the seeds of said plant, so that

25 anthocyanin production in the seeds is visible externally.

34. A process according to claim 33, wherein the genome of said male-sterile parent plant does not contain at least one anthocyanin regulatory gene necessary for the regulation of anthocyanin biosynthesis in the seeds of said plant to produce externally visible anthocyanin in said seeds.

35. The process of claim 34, wherein the genome of said male-sterile parent plant contains a first anthocyanin regulatory gene and the genome of said maintainer parent plant contains a second anthocyanin regulatory gene which, when present with said first anthocyanin regulatory gene in the genome of a plant is capable for conditioning the production of externally visible anthocyanin in seeds.

36. Process to maintain a kit according to claim 33 which comprises:

- crossing said male-sterile parent plant with said maintainer parent plant;
- obtaining the seeds from said male-sterile parent plants and optionally the seeds from said maintainer parent plant in which no anthocyanin is produced; and
- optionally growing said seeds into male-sterile parent plants and maintainer parent plants.

1. Transform a plant of line A with male-sterility gene S
A^{S/-}
2. Transform a plant of line A with color-linked restorer gene Rf:
A^{rf/-}
3. Cross the plants of 1 and 2
Progeny Table of A^{S/-} x A^{rf/-}

	-/-	-/-	-/-
S, -	A ^{S/-} , Rf/-	A ^{S/-} , -/-	A ^{S/-} , -/-
-/-	A ^{-/-} , Rf/-	A ^{-/-} , -/-	A ^{-/-} , -/-

4. Select the progeny from the cross of 3 with both the S and Rf genes (e.g. by means of PCR, or by presence of linked selectable genes)

FIGURE 1

5. Cross the $A^{S/-}, Rf/-$ plants from 4 with $A^{S/-}$.
 Progeny Table of $A^{S/-}, Rf/- \times A^{S/-}, -/-$

	$S, -$	$-/-$
S, Rf	$A^{S/S}, Rf/-$	$A^{S/-}, Rf/-$
$S, -$	$A^{S/S}, -/-$	$A^{S/-}, -/-$
$- , Rf$	$A^{S/-}, Rf/-$	$A^{-/-}, Rf/-$
$-/-$	$A^{S/-}, -/-$	$A^{-/-}, -/-$

6. Select colored seeds, grow into plants and self all plants that contain both the S and Rf genes (e.g. by means of PCR, or by presence of linked selectable marker genes).

Two possible progeny tables depending on the genotype of the selfed plant.

Genotype of selfed plant	Genotype of progeny	Frequency of genotype in progeny	Color phenotype of progeny seeds	Male-fertility phenotype of progeny plants
$S/-, Rf/-$	genotypes as in progeny table in 5	75% 25%	colored seeds non-colored seeds	fertile 75% fertile, 25% sterile
$S/S, Rf/-$	$S/S, Rf/Rf$	25%	colored seeds	fertile
	$S/S, Rf/-$	50%	colored seeds	fertile
	$S/S, -/-$	25%	non-colored seeds	sterile

Progeny which display 75% colored and 25% non-colored seeds and in which all non-colored seeds grow into male-sterile plants are retained.

FIGURE 1 (bis)

3/4

7. Cross the male-sterile plants of the retained progeny of 6 with the male-fertile plants of the retained progeny in 6.
Analyze progeny of each cross. Two types of crosses are possible:
1) $S/S, Rf/Rf \times S/S, -/-$, or
2) $S/S, Rf/- \times S/S, -/-$

Cross	Genotype of progeny	Frequency of genotype in progeny	Color phenotype of progeny seeds	Male-Fertility phenotype of progeny plants
$S/S, Rf/Rf \times S/S, -/-$	$S/S, Rf/-$	100%	colored seeds	fertile
$S/S, Rf/- \times S/S, -/-$	$S/S, Rf/-$	50%	colored seeds	fertile
	$S/S, -/-$	50%	non-colored seeds	sterile

The progeny of the cross which produce 50% colored seeds and 50% non-colored seeds are retained.
The non-colored seedss will all grow into male-sterile first parent plants of this invention, while the colored seeds will all grow into male-fertile second parent plants of this invention.

8. Maintain first and second parent plants by crossing the first and second parent plants and harvesting of the progeny from the male-sterile first parent plant.


Progeny Table of $A^{S/S, -/-} \times A^{S/S, Rf/-}$

	S, Rf	$S, -$
$S, -$	$A^{S/S, Rf/-}$	$A^{S/S, -/-}$

Seeds which will grow into sterile and fertile plants can be separated on the basis of seed color.

FIGURE 1 (ter)

Figure 2





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/29, A01H 1/02, 5/00	A3	(11) International Publication Number: WO 95/34634 (43) International Publication Date: 21 December 1995 (21.12.95)
(21) International Application Number: PCT/EP95/02157 (22) International Filing Date: 6 June 1995 (06.06.95) (30) Priority Data: 08/254,776 6 June 1994 (06.06.94) US (60) Parent Application or Grant (63) Related by Continuation US 08/254,776 (CIP) Filed on 6 June 1994 (06.06.94) (71) Applicant (for all designated States except US): PLANT GENETIC SYSTEMS, N.V. [BE/BE]; Kolonel Bourgstraat 106, B-1040 Brussels (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): KREBBERS, Enno [US/US]; 2014 Branard, Houston, TX 77098 (US). WILLIAMS, Mark [US/BE]; Holdaal 31, B-9000 Gent (BE). LEEMANS, Jan [BE/BE]; Wijngaard 7, B-9831 Deurle (BE). (74) Agents: GUTMANN, Ernest et al.; E. Gutmann - Y. Plasseraud S.A., 3, rue Chauveau-Lagarde, F-75008 Paris (FR).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i> (88) Date of publication of the international search report: 11 January 1996 (11.01.96)
(54) Title: USE OF ANTHOCYANIN GENES TO MAINTAIN MALE STERILE PLANTS		
(57) Abstract <p>A plant consisting essentially of cells which comprise in their genome a homozygous male-sterility genotype at a first genetic locus; and a color-linked restorer genotype at a second genetic locus, which is heterozygous (Rf/-) for a foreign DNA Rf. The foreign DNA Rf comprises: a) a fertility-restorer gene capable of preventing the phenotypic expression of the male-sterility genotype, and b) at least one anthocyanin regulatory gene involved in the regulation of anthocyanin biosynthesis in cells of seeds of the plant which is capable of producing anthocyanin at least in the seeds of the plant, so that anthocyanin production in the seeds is visible externally. Preferably, the anthocyanin regulatory gene is a shortened R, B or C1 gene or a continuation thereof. The invention also relates to DNA sequences encoding shortened R, B or C1 anthocyanin regulatory genes and to a process for maintaining a line of male-sterile plants which comprises crossing a male-sterile parent plant and a maintainer parent plant comprising homozygous male-sterility genotype and a restore genotype comprising fertility-restorer gene and an anthocyanin regulatory gene.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 95/02157

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C12N15/82 C12N15/29 A01H1/02 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MOLECULAR AND GENERAL GENETICS, vol. 242, January 1994 pages 40-48, SCHEFFLER, B., ET AL. 'MOLECULAR ANALYSIS OF C1 ALLELES IN ZEA MAYS DEFINES REGIONS INVOLVED IN THE EXPRESSION OF THIS REGULATORY GENE' see page 42	16
X	THE PLANT JOURNAL, vol. 3, 1993 pages 335-46, CONSONNI, G., ET AL. 'MOLECULAR HOMOLGY AMONG MEMBERS OF THE R GENE FAMILY IN MAIZE' see page 341, right column - page 342 -/--	16

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

10 November 1995

Date of mailing of the international search report

24. 11. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Maddox, A

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>GENES & DEVELOPMENT, vol. 6, no. 5, May 1992, pages 864-875, GOFF, S.A., ET AL. 'FUNCTIONAL ANALYSIS OF THE TRANSCRIPTIONAL ACTIVATOR ENCODED BY THE MAIZE B GENE: EVIDENCE FOR A DIRECT FUNCTIONAL INTERACTION BETWEEN TWO CLASSES OF REGULATORY PROTEINS', see page 865, right column - page 866, left column</p> <p>GENES & DEVELOPMENT, vol. 6, no. 11, November 1992, pages 2152-2164, RADICELLA, J.P., ET AL. 'ALLELIC DIVERSITY OF THE MAIZE B REGULATORY GENE: DIFFERENT LEADER AND PROMOTER SEQUENCES OF TWO B ALLELES DETERMINE DISTINCT TISSUE SPECIFICITIES OF ANTHOCYANIN PRODUCTION', see page 2159, right column see page 2161, right column</p> <p>EMBO JOURNAL, vol. 6, no. 12, 1987 EYNSHAM, OXFORD GB, pages 3553-3558, PAZ-ARES, J., ET AL. 'THE REGULATORY C1 LOCUS OF MAIZE ENCODES A PROTEIN WITH HOMOLOG TO MYB PROTO-ONCOGENE PRODUCTS AND WITH STRUCTURAL SIMILARITIES TO TRANSCRIPTIONAL ACTIVATORS', see the whole document</p>	16
X	<p>GENES & DEVELOPMENT, vol. 6, no. 11, November 1992, pages 2152-2164, RADICELLA, J.P., ET AL. 'ALLELIC DIVERSITY OF THE MAIZE B REGULATORY GENE: DIFFERENT LEADER AND PROMOTER SEQUENCES OF TWO B ALLELES DETERMINE DISTINCT TISSUE SPECIFICITIES OF ANTHOCYANIN PRODUCTION', see page 2159, right column see page 2161, right column</p> <p>EMBO JOURNAL, vol. 6, no. 12, 1987 EYNSHAM, OXFORD GB, pages 3553-3558, PAZ-ARES, J., ET AL. 'THE REGULATORY C1 LOCUS OF MAIZE ENCODES A PROTEIN WITH HOMOLOG TO MYB PROTO-ONCOGENE PRODUCTS AND WITH STRUCTURAL SIMILARITIES TO TRANSCRIPTIONAL ACTIVATORS', see the whole document</p>	16
A	<p>GENES & DEVELOPMENT, vol. 5, no. 2, February 1991, pages 298-309, GOFF, S.A., ET AL. 'IDENTIFICATION OF FUNCTIONAL DOMAINS IN THE MAIZE TRANSCRIPTIONAL ACTIVATOR C1: COMPARISON OF WILD-TYPE AND DOMINANT INHIBITOR PROTEINS', see page 300, right column</p>	16
A	<p>EP, A, 0 412 911 (PLANT GENETIC SYSTEMS NV) 13 February 1991 see page 8, line 40 - line 56</p>	1-36
A	<p>WO, A, 93 25695 (PLANT GENETIC SYSTEMS NV; WILLIAMS MARK (BE); LEEEMANS JAN (BE)) 23 December 1993 see page 15, last paragraph</p>	1-32
A	<p>EP, A, 0 198 288 (ADVANCED GENETIC SCIENCES INC) 22 October 1986 cited in the application see the whole document</p>	1-32

-/--

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
-----------	--	-----------------------

A	US,A,4 727 219 (BRAR GURDIP S ET AL) 23 February 1988 cited in the application see the whole document	1-32
A	WO,A,91 02059 (PIONEER HI BRED INT) 21 February 1991 see the whole document	1-32

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
---	---------------------	----------------------------	---------------------

EP-A-0412911	13-02-91	AU-B-625509 AU-B-6068890 WO-A-9102069 JP-T-3503004	16-07-92 11-03-91 21-02-91 11-07-91
WO-A-9325695	23-12-93	CA-A-2137559 EP-A-0644943 AU-B-5457494 CA-A-2148483 EP-A-0668786 WO-A-9409836	23-12-93 29-03-95 24-05-94 11-05-94 30-08-95 11-05-94
EP-A-0198288	22-10-86	US-A-5180873 AU-B-602306 AU-B-5615586 US-A-5278057	19-01-93 11-10-90 23-10-86 11-01-94
US-A-4727219	23-02-88	NONE	
WO-A-9102059	21-02-91	AU-B-6287390 EP-A-0462231	11-03-91 27-12-91